

UNIVERSITY OF LJUBLJANA
INSTITUTE OF MATHEMATICS, PHYSICS AND MECHANICS
DEPARTMENT OF MATHEMATICS
JADRANSKA 19, 1000 LJUBLJANA, SLOVENIA

Preprint series, Vol. 36 (1998), 598

STRAIGHT-AHEAD WALKS IN
EULERIAN GRAPHS

Tomaž Pisanski Thomas W. Tucker

Arjana Žitnik

ISSN 1318-4865

April 21, 1998

Ljubljana, April 21, 1998

Straight-ahead walks in Eulerian graphs

Tomaž Pisanski, University of Ljubljana, Slovenia

Thomas W. Tucker, Colgate University

Arjana Žitnik, University of Ljubljana, Slovenia

Abstract

A *straight-ahead walk* in an embedded Eulerian graph G always passes from an edge to the opposite edge in the rotation at the same vertex. A straight-ahead walk is called *Eulerian* if all the edges of the embedded graph G are traversed in this way starting from an arbitrary edge. An embedding that contains an Eulerian straight-ahead walk is called an *Eulerian embedding*.

In this article, we characterize some properties of Eulerian embeddings of graphs and of embeddings of graphs such that the corresponding medial graph is Eulerian embedded. We prove that in the case of 4-valent planar graphs, the number of straight ahead walks does not depend on the actual embedding in the plane. Finally, we show that the minimal genus over Eulerian embeddings of a graph can be quite close to the minimal genus over all embeddings.

1 Introduction

Given an Eulerian graph, any matching of edges at each vertex results in a circuit decomposition of the graph. Since there are so many matchings, it would be nice to look at matchings that arise in some natural way or are connected to other properties of the graph. Embeddings of the graph provide an interesting source of matchings. The purpose of this paper is to study the relationship between the embeddings of an Eulerian graph and the circuit decomposition of the graph induced by the embedding by a "straight-ahead" matching. In the other direction, we also show that an Eulerian circuit in a graph can be used to construct interesting embeddings of the graph.

A *straight-ahead walk* or a SAW in the embedded Eulerian graph G always passes from an edge to the opposite edge adjacent to the same vertex; two edges are "opposite" at a vertex of valence $2d$ in an embedded graph if they are d edges apart in the cyclic ordering (*rotation*) of the edges at that vertex induced by the embedding.

In this paper we assume the graphs to be finite and connected and the embeddings to be 2-cell. Let us now introduce some terminology and notation. A *circuit* is a closed walk with no repeated edges. The straight ahead walks, the SAWs, of an embedded Eulerian graph G induce a circuit partition of the edges. Let us denote by $s(G \rightarrow S)$ the number of components of SAW decomposition of G . Notice that this number depends not only on the surface S but also on the given embedding in that surface; it is not hard, for example, to give two embeddings of K_5 in the torus, such that one embedding has two SAWs and the other has three. An embedding of an Eulerian graph G in a surface S is *Eulerian*, if it contains exactly one SAW, ie. $s(G \rightarrow S) = 1$. The *medial graph* of an embedded graph G , $Me(G)$, is a graph, embedded in the same surface as G and is obtained from G as follows: the vertices of $Me(G)$ are the edges of G and two vertices of $Me(G)$ are adjacent if they are adjacent edges in the rotation of a vertex in G . Note that embedded graphs, which are dual to each other, have the same medial graphs. The medial graph of any graph is 4-valent and thus Eulerian. An embedded graph is *Eulerian medial embedded* if its medial is Eulerian embedded.

Eulerian embeddings of 4-valent graphs in the plane are just knot projections (without a specification of which parts of the knot are over or under other parts) and hence are related to Gauss's coding of knot projections (see [7]). An Eulerian embedding of a 4-valent graph in a surface of genus g can be viewed as a knot projection on a genus g Heegard splitting surface for a closed 3-manifold. Unfortunately, the Reidemeister moves for such knot projections include moves across solid handles of the splitting and make knot theory, say for knot polynomials, too complicated. Planar Eulerian graphs are discussed in [4]. Works of Bouchet and others [1, 2, 3, 6, 11] are also related to this paper.

2 Counting SAWs in graphs and medial graphs: some examples

In this section, we give some examples of Eulerian embedded plane graphs and of plane graphs whose medial graph is Eulerian embedded. The most obvious examples of Eulerian embedded graphs are cycles C_n . The medial graphs of odd cycles, which are odd cycles with double edges, are also Eulerian embedded. There exist less trivial infinite families of plane graphs, whose medial graphs are Eulerian embedded, too. It is easy to see, that the medial of the pyramid graph - the antiprism on Figure 1 is Eulerian



Figure 1: The pyramid graph and its medial graph - the antiprism A_4 .

embedded. We used the computer system VEGA, see [8], to verify whether this property holds for all the pyramid graphs. We also checked the number of SAWs in medial graphs of prisms Π_n and antiprisms A_n . The results gave us the following theorem, which we state without proof:

Theorem 1

$$s(A_n \rightarrow Sphere) = \begin{cases} 3 & n = 3k \\ 1 & n \neq 3k \end{cases}$$

$$s(Me(\Pi_n) \rightarrow Sphere) = \begin{cases} 1 & n = 2k + 1 \\ 4 & n = 4k \\ 2 & n = 4k + 2 \end{cases}$$

$$s(Me(A_n) \rightarrow Sphere) = \begin{cases} 4 & n = 3k \\ 2 & n \neq 3k \end{cases}$$

Let G_1 and G_2 be graphs, 2-cell embedded in orientable surfaces S_{k_1} and S_{k_2} , respectively, where S_k denotes the sphere with $k \geq 0$ handles. Let (u_1, v_1) be an edge in G_1 and (u_2, v_2) be an edge in G_2 . If these edges are not both bridges, we can define the *connected sum* $G_1 \# G_2$ of graphs G_1 and G_2 with respect to the directed edges (v_1, u_1) and (v_2, u_2) as follows: take the union of graphs G_1 and G_2 and substitute the edges (v_1, u_1) and (v_2, u_2) by the edges (v_1, v_2) and (u_1, u_2) . The rotation scheme is inherited from the embeddings of G_1 and G_2 , except for the vertices v_1, v_2, u_1 and u_2 . In the rotation around v_1 , u_1 is substituted by v_2 , in the rotation around u_1 , v_1 is substituted by u_2 , and in the rotation around v_2 , u_2 is substituted by v_1 , in the rotation around u_2 , v_2 is substituted by u_1 . The connected sum of G_1 and G_2 is therefore a connected graph, and if at least one of the edges lies

on the boundary of two different faces, the graph $G_1 \# G_2$ is 2-cell embedded in the surface $S_{k_1+k_2}$. The following theorem is very useful for constructing infinite families of Eulerian embedded graphs:

Theorem 2 *Let $G = G_1 \# G_2$. Then $s(G \rightarrow S) = s(G_1 \rightarrow S_{k_1}) + s(G_2 \rightarrow S_{k_2}) - 1$. In particular, if G_1 and G_2 are Eulerian embedded, then G is Eulerian embedded as well.*




Figure 2: The connected sum of the antiprisms A_4 and A_5 .

In Figure 2, the connected sum of the antiprisms A_4 and A_5 is shown. Both A_4 and A_5 are Eulerian embedded and so is their connected sum.

Given an embedded graph, we substitute every k -valent vertex by a cycle on k vertices. The obtained graph is cubic and embedded in the same surface. It is called the *truncation* of the embedded graph. There are two types of faces in a truncated graph: the ones that correspond to former vertices and the ones that correspond to the faces with the boundary twice as long as in the original graph. In [9], the following theorem is proved:

Theorem 3 *The truncations of cubic maps preserve the number of SAWs in their medials.*

So we obtain some other infinite families of Eulerian embedded plane graphs - the medials of all the truncations of the "odd" prisms, medials of their truncations and so on.

3 Number of SAWs in 4-valent plane graphs

Every Eulerian directed graph has an Eulerian embedding, orientable and nonorientable. To obtain such an embedding just choose any embedding

where SAW is the given Eulerian circuit - at each vertex the opposite edges are consecutive in the Eulerian circuit.

But it is not at all obvious how to embed a graph in a given surface with the minimal possible number of SAWs or to find the surface of minimal genus in which a graph G can be embedded so to have only one SAW. These questions seem to be very difficult and are still open. Nevertheless, for the plane the following result holds:

Theorem 4 *Let G be a planar 4-valent graph. Then the number of SAWs is the same for any embedding of G in the plane.*

Proof For 3-connected graphs the theorem trivially holds, since they have essentially unique embeddings in the plane.

For 2-connected graphs the proof depends on the well-known theorem, that any embedding of a planar 2-connected graph can be obtained from another by a sequence of operations dual to the Whitney's 2-switchings. This operation is defined as follows: if we have a separation pair $\{x, y\}$, we turn around one component of a graph, adjacent to x and y ; so the orders of neighbors of x and y in this component are reversed. This procedure is illustrated in Figure 3.




Figure 3: An example of a dual 2-switching

The proof consists of considering of all possible cases of how SAWs can pass through a separation pair. As an example, let us consider the case, where there is only one SAW passing through x and y , and it passes first twice through x and then twice through y . After the dual 2-switching, the SAW through x and y is changed, but the number of SAWs in G remains the same, see Figure 3, where the SAWs through x and y are depicted in bold lines and the rest of the graph, in which the dual 2-switching doesn't affect the SAWs, is depicted in gray.

If G is not 2-connected, it has a cut-vertex, say v . Through the cut-vertex v , only one SAW can pass. Changing the rotation at v such that the

embedding remains plane does not change the number of SAWs through v . \square

This theorem does not hold for all planar Eulerian graphs. In Figure 4 two embeddings in the plane of the same graph are shown, which contain different numbers of SAWs.



Figure 4: An example of a planar graph having different number of SAWs in different embeddings in the plane.

But from the proof of the Theorem 4 it can easily be seen that the Theorem holds for a more general class of 4-valent graphs, namely the planar Eulerian graphs with cut-vertices and separation pairs of degree not different from 4.

Corollary 5 *Let G be a planar Eulerian graph with possible cut-vertices and separation pairs of degree 4. Then the number of SAWs is independent of the embedding of G in the plane.*

4 Eulerian medial embeddings

Any 2-cell embedding of a connected graph G can be represented by a triple (G, P, λ) , where P is the *rotation scheme* of G and $\lambda : E(G) \rightarrow \{-1, 1\}$ assigns *signatures* to the edges, which tells us, whether an edge is orientation preserving or orientation reversing, see [12].

Given an embedding of a graph G , we change the signatures of the edges such that the orientation preserving edges become orientation reversing and vice versa. A different embedding of G is obtained, which is called the *Petrie dual* of (the embedded) graph G . The faces of the Petrie dual are called *Petrie walks* of the original embedding of G . It is not hard to see that

SAWs of medial graphs correspond to Petrie walks of the original map. See, for example, [7], where the Petrie walks are called left-right paths. That means, that an Eulerian medial embedding of a graph is equivalent to Petrie dual being 1-face embedded.

Theorem 6 *Every graph embedding can be subdivided to give an Eulerian medial embedding.*

Proof The proof depends on the following idea: If SAWs of a 4-valent graph have two circuits at a vertex the other two matchings at a vertex give one circuit through that vertex. Subdividing an edge of the original graph can be viewed as changing the matching of the corresponding vertex of the medial graph. At each step we subdivide an edge, whose corresponding vertex of the medial graph is contained in two different SAWs, and at the end we obtain an Eulerian medial embedded graph. \square

The following corollary is an easy consequence of the Theorem and the fact that for every surface there exist medial graphs.

Corollary 7 *Every surface admits Eulerian embeddings.*

The question arises, whether every graph has an Eulerian medial embedding. If we consider only orientable surfaces, the answer is "no". The simplest example of graphs having no orientable Eulerian embedding are even cycles. The embedding of an even cycle to an orientable surface is unique and the corresponding medial graph has two SAWs. Let us define a *cactus* as a graph, in which every vertex belongs to at most one cycle.

Theorem 8 *In a cactus, the number of SAWs in the medial is equal to the number of even cycles + 1.*

Proof By induction. \square

Note, that Theorem 8 is not valid for a similar class of graphs with the property that each *edge* belongs to at most one cycle.

Attaching a graph G_1 to graph G_2 by an edge is the following procedure: we choose edges e_1 in G_1 and e_2 in G_2 , subdivide e_i and denote the additional vertex by v_i , $i = 1, 2$. Then we join the vertices v_1 and v_2 by an edge.

Corollary 9 *If a cactus with even cycles is attached by an edge to an arbitrary graph G , then the resulting graph doesn't have an Eulerian medial embedding.*

These examples of graphs are not even 2-connected. The graph of a 3-dimensional cube, usually denoted by Q_3 , is a 3-connected cubic graph. It has 2^8 different embeddings (many of them are equivalent). We have counted the numbers of SAWs in the medials of all these embeddings of Q_3 with the help of a computer and found out, that they always have more than one SAW. The question arises, which 3-connected graphs do have an Eulerian medial embedding. In particular, is it true that a graph with a 1-face embedding has an Eulerian medial embedding?

If we also allow nonorientable embeddings, every graph has an Eulerian medial embedding.

Theorem 10 *For every rotation scheme, there is an assignment of signatures to edges that gives an Eulerian medial embedding (possibly nonorientable).*

Proof The proof is divided in two steps.

- Change the signatures of edges between distinct faces until a one-face embedding is obtained. If the signature of an edge between two faces is changed, these two faces are merged to one face.
- The Petrie dual of the so-obtained graph has the medial with required property.

□

5 Bounds on Eulerian genus

Every Eulerian directed graph has an Eulerian embedding, orientable and non orientable. To obtain such an embedding just choose any embedding where the SAW is the given Eulerian circuit - at each vertex the opposite edges are consecutive in the Eulerian circuit. We can define the *Eulerian genus* of a graph G as the smallest possible genus of an orientable surface, in which G can be Eulerian embedded. In section 2, we have seen some examples of planar graphs which are Eulerian embedded in the plane. In Figure 5 the embedding of K_5 in the torus is shown. It only has one SAW, which means, that the Eulerian genus of K_5 is equal to its ordinary genus.

Lemma 11 *Let G be an Eulerian graph, embedded in a surface of genus g with $s(G \rightarrow S_g) = k$. Then the Eulerian genus of G is less or equal to $g + k - 1$.*




Figure 5: An Eulerian embedding of K_5 in the torus.

Proof Let e and f be two edges, adjacent in the rotation at a vertex v , and let them belong to different SAWs (if there is more than 1 SAW, this must happen). Switching e and f at v causes the SAWs through e and f to be joined into one SAW. We repeat this procedure until there is only one SAW left. Switching the rotation at a vertex can only increase the genus by one (see, for example [5]). So after $k - 1$ switches, the genus is increased by at most $k - 1$. \square

Remark: let $(\dots, e_1, e_2, \dots, e_k, \dots)$ be the rotation at a vertex v and let the edges e_1, \dots, e_k belong to distinct SAWs. Then changing the rotation at v to $(\dots, e_2, \dots, e_k, e_1, \dots)$ causes all these SAWs to join.

Corollary 12 *The Eulerian genus of $C_m \times C_n$ is less or equal to $m + n$.*

Proof The graph $C_m \times C_n$ can be embedded in the torus in the obvious way such that it contains $n + m$ SAWs. It follows from the Lemma, that the Eulerian genus must be at most $1 + (m + n - 1) = m + n$. \square

Let us state a theorem, characterizing the number of SAWs in covering graphs. For the definitions of covering graphs and Cayley graphs see, for example, [5].

Theorem 13 *Let G be an embedded voltage graph with voltages from group α of order n . Let the SAWs of G be C_1, C_2, \dots, C_k and let the product of voltages along C_i have order m_i in the voltage group α , $i = 1, 2, \dots, k$. (The voltage on a minus directed edge is understood to be the group inverse of the voltage on its reverse edge.) Then the derived graph \tilde{G} has $\frac{n}{m_1} + \frac{n}{m_2} + \dots + \frac{n}{m_k}$ SAWs.*

Proof The proof is based on the following theorem from [5]: Let C be a k -cycle in the base space of an ordinary voltage graph (G, α) such that the

product of voltages along C has order m in the voltage group α . Then each component of the preimage $p^{-1}(C)$ is a $k \cdot m$ -cycle and there are $|\alpha|/m$ such components. \square

Corollary 14 *Let G be an Eulerian embedded graph, which is Eulerian directed according to its SAW. Given any cyclic voltage graph on G such that the product of voltages along the directed edges generates the group, then the covering graph is Eulerian embedded.*

Cayley graphs are regular coverings of bouquets of circles. A regular embedding of a Cayley graph is given by lifting the rotation of the bouquet of circles to the Cayley graph. The rotation is called *special*, if the SAW in the bouquet of circles is Eulerian. The following Corollary is thus an easy consequence of the Corollary 14.

Corollary 15 *Given any regular embedding of an even Cayley graph, it is Eulerian if and only if the group is cyclic, the rotation is special and the product of the generators along the SAW in the bouquet of circles generates the group.*




Figure 6: A triangular embedding of B_9 in the double torus.

Example 1 *Figure 6 shows the embedding of the bouquet of 9 circles B_9 in the double torus. Its edges are directed and have voltages from the group Z_{19} . The covering graph of this voltage graph is a triangular embedding of K_{19} and thus a minimal genus embedding.*

The number of SAWs in the embedding of B_9 is 2, the products of voltages along the SAWs are 10 and 3 in Z_{19} , and are relatively prime to 19. By the theorem 13, the triangular embedding of K_{19} , obtained from (B_9, Z_{19}) , contains two SAWs and we conclude that the Eulerian genus of K_{19} differs from its ordinary genus by at most 1.

The construction can be generalized to all complete graphs on $12t + 7$ vertices, which was done by G. Ringel in 1961 (see [10] or [5]). With help of a computer we have constructed the graphs B_{6t+3} , which give us the triangular embeddings of K_{12t+7} as covering graphs. We calculated the numbers of SAWs in B_{6t+3} , $t = 1, \dots, 150$, and the products of voltages along the SAWs. Part of the results is given in the tables 1 and 2. It is interesting, that there exist also large t such that the Eulerian genus of K_{12t+7} differs from the ordinary genus of K_{12t+7} by at most 1.

t=	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
$s(B_{6t+3} \rightarrow S_{t+1})$	2	3	6	3	2	7	4	5	6	7	2	9	6	3	14	3

t=	17	18	19	20	21	22	23	24	25	26	27	28	29	30
$s(B_{6t+3} \rightarrow S_{t+1})$	2	9	2	3	6	9	6	15	2	3	6	5	8	9

Table 1: Numbers of SAWs in the graphs B_{6t+3} .

t=	Group	Products of voltages along the SAWs									
1	Z_{19}	10	3								
2	Z_{31}	20	18	5							
3	Z_{43}	17	20	6	12	7	20				
4	Z_{55}	13	0	9							
5	Z_{67}	12	11								
6	Z_{79}	38	57	20	2	22	13	37			
7	Z_{91}	76	15	1	15						
8	Z_{103}	8	42	50	63	17					
9	Z_{115}	20	35	20	32	19	54				
10	Z_{127}	63	110	46	48	95	112	21			
11	Z_{139}	94	23								
12	Z_{151}	121	132	22	97	29	128	42	25	71	

Table 2: Products of voltages along the SAWs in the graphs B_{6t+3} .

6 Conclusion and open problems

The natural question is which Eulerian graphs have their Eulerian genus equal to the ordinary genus. Another question that can be posed is the

following: which 2-cell embeddings of graphs have their connected and four-valent medial graphs Eulerian embedded? Finally, which graphs have at least one orientable embedding such that the corresponding medial graph is Eulerian embedded?

References

- [1] L.D. Andersen, A. Bouchet and B. Jackson, A-trails in 4-regular graphs, *J. Combin. Theory Ser. B* 66 (1996), 233-246.
- [2] A. Bouchet, Multimatroids II. Orthogonality, minors and connectivity, *The Electronic Journal of Combinatorics*, Volume 5(1) R8.
- [3] A. Bouchet, Compatible Euler tours and supplementary Eulerian vectors, *Europ. J. Combin.*, 514 (1993), 513-520.
- [4] H. Fleischner, *Eulerian Graphs and Related Topics*, Part I, Vols. 1 and 2, North Holland, Amsterdam, 1990-1991.
- [5] J.L. Gross, & T.W. Tucker, *Topological Graph Theory*, J. Wiley & sons, 1987.
- [6] D.-Y. Jeong, Realizations with a cut-through Eulerian circuit, *Discrete Mathematics* 137 (1995) 265-275.
- [7] S. Lins, B. Richter & H. Shank, The Gauss code problem off the plane, *Aequationes Mathematicae*, 33 (1987) 81-95.
- [8] T. Pisanski & coworkers, VEGA, a programming tool for manipulating discrete mathematical structures, see <http://vega.ijp.si/Html/doc/Vega03.html>.
- [9] T. Pisanski, T.W. Tucker & A. Žitnik, Eulerian embeddings of graphs, submitted.
- [10] G. Ringel, *Map Color Theorem*, Springer - Verlag, Berlin, 1984.
- [11] H. Shank, The theory of left-right paths, in: *Combinatorial Mathematics III. Lecture Notes in Math.* 452. Springer, Berlin, 1975, pp. 42-54.
- [12] S. Stahl, Generalized embedding schemes, *J. Graph Theory*, Vol. 2 (1978) 41-52.