

UNIVERSITY OF LJUBLJANA
INSTITUTE OF MATHEMATICS, PHYSICS AND MECHANICS
DEPARTMENT OF MATHEMATICS
JADRANSKA 19, 1000 LJUBLJANA, SLOVENIA

Preprint series, Vol. 39 (2001), 779

COMPUTING GRAPH
INVARIANTS ON ROTAGRAPHS
USING DYNAMIC ALGORITHM
APPROACH: THE CASE OF
(2, 1)-COLORINGS AND
INDEPENDENCE NUMBERS

Sandi Klavžar Aleksander Vesel

ISSN 1318-4865

October 22, 2001

Ljubljana, October 22, 2001

Computing graph invariants on rotagraphs using dynamic algorithm approach: the case of (2,1)-colorings and independence numbers

Sandi Klavžar*

Department of Mathematics, PEF, University of Maribor
Koroška cesta 160, 2000 Maribor, Slovenia
sandi.klavzar@uni-lj.si

Aleksander Vesel*

Department of Mathematics, PEF, University of Maribor
Koroška cesta 160, 2000 Maribor, Slovenia
vesel@uni-mb.si

October 6, 2001

Abstract

Rotagraphs generalize all standard products of graphs in which one factor is a cycle. A computer based approach for searching graph invariants on rotagraphs is proposed and two of its applications are presented. First, the λ -numbers of the Cartesian product of a cycle and a path are computed, where the λ -number of a graph G is the minimum number of colors needed in a (2,1)-coloring of G . The independence numbers of the family of the strong product graphs $C_7 \boxtimes C_7 \boxtimes C_{2k+1}$ are also obtained.

Key words: rotagraph, dynamic algorithm, (2,1)-coloring, independence number, Cartesian product of graphs, strong product of graphs

AMS subject classification: 05C85, 05C15, 05C69, 68R10

Proposed running head: (2,1)-colorings and independence numbers of rotagraphs

*Supported by the Ministry of Education, Science and Sport of Slovenia under the grant 0101-504.

1 Introduction

A large part of Graph Theory consists of the study of graph invariants, as for instance the chromatic number, the independence number, and the domination number. Unfortunately, the determination of each of these invariants is NP-hard. Therefore many approximation and heuristic algorithms have been developed which obtain near-optimal solutions within reasonable running times. However, such algorithms in general provide only upper and lower bounds.

One way to approach such invariants is to decompose a graph into smaller and simpler constituents by some basic operations and try to gain a global information from the constituents. For instance, one might try to decompose a graph with respect to one of the standard graphs products, see [15]. Quite often, product graphs having cycles as factors are important—as is the case of the Shannon capacity. A natural generalization of such structures is formed by rotagraphs that are in turn an important class of polygraphs. The later were introduced in chemical graph theory as a model for polymers, cf. [1], and studied in, for instance, [19, 20, 30].

In this paper we propose a dynamic algorithm approach for computing graph properties of rotagraphs. The main idea is to build a function (an invariant) on a rotagraph from corresponding functions on its basic building blocks. Then the problem reduces to a search for a certain subgraph in an associated directed graph. In particular, if the function in question is rather local, one has to look for directed cycles. Although finding a cycle of length n , for an arbitrary n , is NP-complete, we were able to apply our approach for two central invariants. The main reason for the success is that it suffices to deal with small strongly connected components, therefore the computations can be done in a reasonable time.

We first succeeded to compute the λ -numbers of the Cartesian product of a cycle and a path, for which partial results were obtained in [18]. The λ -number of a graph G is the minimum number of colors needed in a (2,1)-coloring of G . The concept of $L(2,1)$ -coloring (or (2,1)-labeling) was introduced by Griggs and Yeh in [12], and has since been extensively studied, see for instance [3, 4, 6, 14, 21, 22]. In particular, the λ -numbers (and their generalizations) of Cartesian product graphs (like hypercubes and Hamming graphs) were treated in [5, 7, 8, 12, 18, 29]. In addition, Jha obtained the λ -numbers of some infinite families of Cartesian products of several cycles [16] as well as of strong products of several cycles [17].

In the second application we determine the independence numbers of the strong products $C_7 \boxtimes C_7 \boxtimes C_{2k+1}$. Studies of the independence number of the strong product of odd cycles have been inspired by the Shannon's work [23] on the determination of the zero-capacity of a noisy channel. Shannon formulated the problem in the form of graph theory and supplied some partial results. It turns out that the solution of the problem requires the determination of the independence number of the strong product of graphs containing odd cycles as factors. Note also that the Shannon capacity of the 7-cycle is still open [11, 27].

2 Definitions

Let $G = (V(G), E(G))$ be a graph. A *walk* is a sequence of vertices v_1, v_2, \dots, v_n and edges $v_i v_{i+1}$, $1 \leq i \leq n-1$. A *path* on n vertices is a walk on n different vertices and denoted P_n . A walk is *closed* if $v_1 = v_n$. A closed walk in which all vertices (except the first and the last) are different, is a *cycle*. The cycle on n vertices is denoted C_n . For $u, v \in V(G)$, $d_G(u, v)$ or $d(u, v)$ denotes the length of a shortest walk (i.e., the number of edges on a shortest walk) in G from u to v . These definitions extend naturally to directed graphs.

The *Cartesian product* of graphs G and H is the graph $G \square H$ with vertex set $G \times H$ and $(x_1, x_2)(y_1, y_2) \in E(G \square H)$ whenever $x_1 y_1 \in E(G)$ and $x_2 = y_2$, or $x_2 y_2 \in E(H)$ and $x_1 = y_1$. The *strong product* of graphs G and H is the graph $G \boxtimes H$ with vertex set $G \times H$ and $(x_1, x_2)(y_1, y_2) \in E(G \boxtimes H)$ whenever $x_1 y_1 \in E(G)$ and $x_2 = y_2$, or $x_2 y_2 \in E(H)$ and $x_1 = y_1$, or $x_1 y_1 \in E(G)$ and $x_2 y_2 \in E(H)$. The Cartesian product and the strong product are commutative and associative, having the trivial graph as a unit, cf. [15].

A $L(2, 1)$ -coloring of a graph G is a function f from the vertex set $V(G)$ to the set of non-negative integers C (called *colors*) such that

$$\begin{aligned} |f(u) - f(v)| &\geq 2 \text{ if } d(u, v) = 1, \text{ and} \\ |f(u) - f(v)| &\geq 1 \text{ if } d(u, v) = 2. \end{aligned}$$

An k - $L(2, 1)$ -coloring is a $L(2, 1)$ -coloring of G such that $C = \{0, \dots, k\}$. An optimal $L(2, 1)$ -coloring of G is a k - $L(2, 1)$ -coloring with k smallest possible. The largest color used by an optimal $L(2, 1)$ -coloring is called the λ -number of G and denoted by $\lambda(G)$. Note that the overall numbers of colors involved in an optimal coloring is $\lambda(G) + 1$. Recall that the problem of determining the λ -number is NP-hard [9].

A set $S \subseteq V(G)$ is *independent* if $xy \notin E(G)$ for any pair of vertices $x, y \in S$. Cardinality of a largest independent set S of G is the *independence number* of G and denoted $\alpha(G)$.

Let G_0, G_1, \dots, G_{n-1} be disjoint graphs and X_0, X_1, \dots, X_{n-1} a sequence of sets of edges such that an edge of X_i joins a vertex of G_i with a vertex of G_{i+1} (indices modulo n). A *polygraph*

$$\Omega_n = \Omega_n(G_0, G_1, \dots, G_{n-1}; X_0, X_1, \dots, X_{n-1})$$

is defined in the following way:

$$\begin{aligned} V(\Omega_n) &= V(G_0) \cup V(G_1) \cup \dots \cup V(G_{n-1}), \\ E(\Omega_n) &= E(G_0) \cup X_0 \cup E(G_1) \cup X_1 \cup \dots \cup E(G_{n-1}) \cup X_{n-1}. \end{aligned}$$

Assume that for $0 \leq i \leq n-1$, G_i is isomorphic to a fixed graph G . Let, in addition, the sets X_i , $0 \leq i \leq n-1$, be equal to a fixed edge set X . Then we call the polygraph Ω_n a *rotagraph* and denote it $\omega_n(G; X)$. We will also say that $\omega_n(G; X)$ is a rotagraph with consecutive *fibers* G_0, G_1, \dots, G_{n-1} . A *fasciagraph* $\psi_n(G; X)$ is a rotagraph $\omega_n(G; X)$ without edges between the fibers G_{n-1} and G_0 . An example of a rotagraph is given in Fig. 1. Its fibers (one being encircled in the figure) are isomorphic to $K_2 \cup K_1$.




Figure 1: Rotagraph and its fiber

3 Graph properties and rotagraphs

For a graph G set

$$\mathcal{F}_q(G) = \{f : V(G) \rightarrow \{0, 1, \dots, q-1\}\}.$$

A subset of $\mathcal{F}_q(G)$ will be called a *graph q -property*. If q will be clear from the context or not essential, we will shortly say a *graph property*. Let us first list some examples of graph properties.

1. Let $\mathcal{C}_q(G) \subseteq \mathcal{F}_q(G)$ be the set of functions f with the following characterizing property: If $uv \in E(G)$ then $f(u) \neq f(v)$. It is clear that $\mathcal{C}_q(G)$ represents the set of all proper q -colorings of G . In particular $\mathcal{C}_q(G) \neq \emptyset$ if and only if $\chi(G) \leq q$.
2. Let $\mathcal{I}_2(G) \subseteq \mathcal{F}_2(G)$ be the set of functions f with the following property: Let $f \in \mathcal{I}_2(G)$, then there is an independent set I of G such that $I = f^{-1}(1)$. Then $\mathcal{I}_2(G)$ is the set of characteristic functions of the independent sets of G .
3. Let the functions f of $\mathcal{D}_2(G) \subseteq \mathcal{F}_2(G)$ be defined with the following property: For any $f \in \mathcal{D}_2(G)$ and any vertex $u \in V(G)$ with $f(u) = 0$ there exists a vertex $v \in N(u)$ such that $f(v) = 1$. Observe that $\mathcal{D}_2(G)$ represents the dominating sets of G .
4. Let $\mathcal{L}_q(G) \subseteq \mathcal{F}_q(G)$ be the set of functions f with the following property: Let $f \in \mathcal{L}_q(G)$, then if $uv \in E(G)$ we have $|f(u) - f(v)| \geq 2$, and if $d(u, v) = 2$ we have $|f(u) - f(v)| \geq 1$. Clearly, $\mathcal{L}_q(G)$ describes the admissible $(2, 1)$ -colorings of G .
5. Let $\mathcal{B}_2(G) \subseteq \mathcal{F}_2(G)$ be the set of functions f with the following property: Let $f \in \mathcal{B}_2(G)$, then $f^{-1}(0)$ and $f^{-1}(1)$ are independent sets of G . Then $\mathcal{B}_2(G)$ describes the bipartitions of G . In particular $\mathcal{B}_2(G) \neq \emptyset$ if and only if G is bipartite.

We now turn our attention to rotagraphs. Let $\omega_n(G; X)$ be a rotagraph with consecutive fibers G_0, G_1, \dots, G_{n-1} . Then the restriction of $f \in \mathcal{F}_q(\omega_n(G; X))$ to consecutive fibers $X_i, X_{i+1}, \dots, X_{i+k}$ (indices modulo n) will be denoted f_i^{i+k} . We say that a graph

property \mathcal{P}_q is *hereditary* (for rotagraphs), if for any rotagraph $\omega_n(G; X)$ with consecutive fibers G_0, G_1, \dots, G_{n-1} ,

$$f \in \mathcal{P}_q(\omega_n(G; X)) \Rightarrow f_i^{i+k} \in \mathcal{P}_q(\Psi_{k+1}(G; X)); \quad i, k = 0, 1, \dots, n-1.$$

Note that $\mathcal{C}_q, \mathcal{I}_2, \mathcal{L}_q$, and \mathcal{B}_2 are hereditary properties, while \mathcal{D}_2 is not. To see that \mathcal{D}_2 is not hereditary, consider the rotagraph of Fig. 1 and its dominating set consisting of the middle vertices of each fiber.

A graph property \mathcal{P}_q is called *d-local* (for rotagraphs), $d \geq 1$, if for any rotagraph $\omega_n(G; X)$, $n \geq 2d+1$, with consecutive fibers G_0, G_1, \dots, G_{n-1} , and any $f \in \mathcal{F}_q(\omega_n(G; X))$,

$$f_i^{i+d} \in \mathcal{P}_q(\Psi_{d+1}(G; X)), 0 \leq i \leq n-1 \Rightarrow f \in \mathcal{P}_q(\omega_n(G; X)).$$

Note that $\mathcal{C}_q, \mathcal{I}_2$, and \mathcal{D}_2 are 1-local properties (for rotagraphs), \mathcal{L}_q is a 2-local property, while \mathcal{B}_2 is not *d-local* for any $d \geq 1$.

The basic building stone of a rotagraph $\omega_n(G; X)$ is the fasciagraph $\Psi_2(G; X)$. Therefore, our main idea is to build functions (alias graph properties) on $\omega_n(G; X)$ from corresponding functions on $\Psi_2(G; X)$. We proceed as follows.

Let \mathcal{P}_q be a *d-local* property, and $\omega_n(G; X)$ a rotagraph with $n \geq 2d+1$. We define a directed graph $D_d(G; X)$ as follows. Its vertices are the functions from $\mathcal{P}_q(\Psi_2(G; X))$. Then make an arc from f to g if and only if f restricted to the second fiber of $\Psi_2(G; X)$ equals to g restricted to the first fiber of $\Psi_2(G; X)$. In addition, if $d \geq 2$, then for any directed path of length $d-1$, say $f_1 \rightarrow f_2 \rightarrow \dots \rightarrow f_d$, we make an arc from f_1 to f_d if the composition of f_1, f_2, \dots, f_d belongs to $\mathcal{P}_q(\Psi_{d+1}(G; X))$. In the particular case when $d=2$ we interpret this as follows: If the composition of f_1 and f_2 belongs to $\mathcal{P}_q(\Psi_3(G; X))$ then we leave the arc from f_1 to f_2 , otherwise we remove it. Now we can state:

Theorem 1 *Let \mathcal{P}_q be a hereditary, d-local property, and $\omega_n(G; X)$ a rotagraph with $n \geq 2d+1$. Then $\mathcal{P}_q(\omega_n(G; X)) \neq \emptyset$ if and only if $D_d(G; X)$ contains (not necessarily different) vertices f_0, f_1, \dots, f_{n-1} connected with arcs (f_i, f_{i+1}) and (f_i, f_{i+d-1}) for $i = 0, 1, \dots, n-1$ (indices modulo n).*

Proof. Let G_0, G_1, \dots, G_{n-1} be the consecutive fibers of $\omega_n(G; X)$.

Assume $\mathcal{P}_q(\omega_n(G; X)) \neq \emptyset$ and let $f \in \mathcal{P}_q(\omega_n(G; X))$. Since \mathcal{P}_q is hereditary, f_i^{i+1} is a vertex of $D_d(G; X)$ for $i = 0, 1, \dots, n-1$. Set $f_i = f_i^{i+1}$, $i = 0, 1, \dots, n-1$. Clearly, there is an arc from f_i to f_{i+1} for all i . If $d=1$ we are done, so let $d \geq 2$. Then, again as \mathcal{P}_q is hereditary, f_i^{i+d-1} belongs to $\mathcal{P}_q(\Psi_{d+1}(G; X))$ and we have an arc from f_i to f_{i+d-1} .

Suppose now that $D_d(G; X)$ contains vertices f_0, f_1, \dots, f_{n-1} and arcs (f_i, f_{i+1}) and (f_i, f_{i+d-1}) . Define a function $f \in \mathcal{F}_q(\omega_n(G; X))$ with the property that f_i^i is equal to f_i restricted to the first fiber of $\Psi_2(G; X)$. Since we have an arc from f_i to f_{i+d-1} , f_i^{i+d-1} belongs to $\mathcal{P}_q(\Psi_{d+1}(G; X))$. Moreover, as \mathcal{P}_q is a *d-local* property we conclude that $f \in \mathcal{P}_q(\omega_n(G; X))$. \square

The most practically important special case of the theorem is:

Corollary 2 *Let \mathcal{P}_q be a hereditary, d -local property, $1 \leq d \leq 2$, and $\omega_n(G; X)$ a rotagraph with $n \geq 5$. Then $\mathcal{P}_q(\omega_n(G; X)) \neq \emptyset$ if and only if $D_d(G; X)$ contains a directed closed walk of length n .*

4 $L(2, 1)$ -colorings of the Cartesian product of a cycle and a path

In this section we apply the proposed general approach from Section 3 to obtain the λ -numbers of the Cartesian product of a path and a cycle. Whittlesey, Georges, and Mauro [29] obtained the λ -numbers of the Cartesian product of two paths. They proved that if $m \geq 4$, then $\lambda(P_m \square P_2) = 5$ and if $m, n \geq 4$, or $m \geq 5$ and $n \geq 3$, then $\lambda(P_m \square P_n) = 6$. For the Cartesian product of paths and cycles, the following partial results from [18] are known:

Theorem 3 *Let $m \geq 3$. Then:*

- (i) *If $m \equiv 0 \pmod{3}$, then $\lambda(C_m \square P_2) = 5$.*
- (ii) *If $m \not\equiv 0 \pmod{3}$, then $\lambda(C_m \square P_2) \leq 6$.*
- (iii) *If $m \equiv 0 \pmod{7}$ and $n \geq 3$, then $\lambda(C_m \square P_n) = 6$.*
- (iv) *If $m, n \geq 3$, then $\lambda(C_m \square P_n) \leq 7$.*

In this section we show:

Theorem 4 *Let $m \geq 3$. Then:*

- (i) $\lambda(C_m \square P_2) = \begin{cases} 5; & m \equiv 0 \pmod{3}, \\ 6; & \text{otherwise.} \end{cases}$
- (ii) $\lambda(C_m \square P_3) = \begin{cases} 7; & m = 4, 5, \\ 6; & \text{otherwise.} \end{cases}$
- (iii) For $n \geq 4$, $\lambda(C_m \square P_n) = \begin{cases} 6; & m \equiv 0 \pmod{7}, \\ 7; & \text{otherwise.} \end{cases}$

In the rest of this section we prove Theorem 4. For this sake we need the following lower bound due to Griggs and Yeh [12]:

Lemma 5 *Let G be graph with maximum degree $\Delta \geq 2$. If G contains three vertices of degree Δ such that one of them is adjacent to the other two, then $\lambda(G) \geq \Delta + 2$.*

Case (i): $\lambda(C_m \square P_2)$.

The Cartesian product $C_m \square P_2$ is isomorphic to the rotagraph $\omega_m(P_2; X)$, where X consists of two edges inducing an identity between the corresponding fibers, that is, between the P_2 's. Consider the graph property \mathcal{L}_6 and recall that it is hereditary, 2-local property. From Corollary 2 we know that $\mathcal{L}_6(\omega_m(P_2; X)) \neq \emptyset$ if and only if $D_2(P_2; X)$ contains




Figure 2: Graph $D_2(P_2; X)$ (without isolated vertices)

a directed closed walk of length m . The graph $D_2(P_2; X)$ (without isolated vertices) is depicted in Fig. 2.

Fig. 2 immediately implies that all directed closed walks in $D_2(P_2; X)$ are of length $3k$, $k \geq 1$. By Lemma 5, $\lambda(C_m \square P_2) \geq 5$, and by Theorem 3, $\lambda(C_m \square P_2) \leq 6$, hence the conclusion.

Case (ii): $\lambda(C_m \square P_3)$.

For $m = 4, 5$ the result follows from a simple computer program based on backtracking. The same program gives the 6- $L(2, 1)$ -colorings for $m = 3k$ (Fig. 3), for $m = 3k + 1$ (Fig. 4), and for $m = 3k + 2$ (Fig. 5).

Since by Lemma 5, $\lambda(C_m \square P_3) \geq 6$, Case (ii) is settled.

Case (iii): $\lambda(C_m \square P_n)$, $n \geq 4$.

For $m \equiv 0 \pmod{7}$ the conclusion follows from Theorem 3.

Let $n = 4$ and consider the rotagraph $\omega_m(P_4; X)$, where X consists of four edges inducing an identity between the corresponding P_4 's. Consider the graph property \mathcal{L}_7 . Corollary 2 implies that $\omega_m(P_4; X)$ has a 6- $L(2, 1)$ -coloring if and only if $D_2(P_4; X)$ contains a directed closed walk of length m . The graph $D_2(P_4; X)$ consists of 1248 vertices and 912




Figure 3: 6- $L(2, 1)$ -coloring of $C_{3k} \square P_3$, $k \geq 1$




Figure 4: 6- $L(2, 1)$ -coloring of $C_{3k+1} \square P_3$, $k \geq 2$

edges (determined by a computer program). In order to search for cycles in $D_2(P_4; X)$, the strongly connected components were detected first. Linear-time algorithms that obtain strongly connected components are well known (see e.g. [28]). Eight strongly connected components of $D_2(P_4; X)$ were computed, each of them consisting of seven vertices and exactly one directed cycle. Therefore, all directed closed walks in H are of length $7k$, $k \geq 1$, thus a 6- $L(2, 1)$ -coloring of $\omega_m(P_4; X)$ for $m \not\equiv 0 \pmod{7}$ does not exist. Since $\lambda(C_m \square P_4) \leq 7$ by Theorem 3, and $\lambda(C_m \square P_4) \geq 6$ by Lemma 5, the argument for the subcase $n = 4$ of is complete.

For $n \geq 5$ just note that $C_m \square P_4$ is a subgraph of $C_m \square P_n$, $n \geq 4$, and apply Lemma 5 and Theorem 3.




Figure 5: 6- $L(2, 1)$ -coloring of $C_{3k+2} \square P_3$, $k \geq 2$

5 Independent sets in strong products of cycles

The independence numbers of the strong product of cycles where at most one cycle is odd are well-known. The problem is incomparably more difficult for the strong product of cycles with at least two odd factors. Hales [13], as well as Sonnemann and Krafft [24], proved that for $j, k \in \mathbb{N}$, $j \geq k$, $\alpha(C_{2j+1} \boxtimes C_{2k+1}) = jk + \lfloor k/2 \rfloor$. For the strong product of three (or more) odd cycles only partial results are known. In [25], see also [2], the independence numbers of the strong products of three cycles of moderate sizes were considered: $C_5 \boxtimes C_7 \boxtimes C_{2k+1}$, $C_5 \boxtimes C_9 \boxtimes C_{2k+1}$, $C_7 \boxtimes C_7 \boxtimes C_{2k+1}$, and $C_7 \boxtimes C_9 \boxtimes C_{2k+1}$. The results are summarized in the following table.

cycle lengths	k	α
5, 7, k	$5 \geq k \geq 3$	$7k + 2$
5, 7, k	$k \geq 6$	$7k + 3$
5, 9, k	$k \geq 4$	$9k + 4$
7, 7, 7		33
7, 7, 9		44
7, 7, k	$k \geq 5$	$10k + 4$ or $10k + 5$
7, 9, k	$k \geq 4$	$13k + 5$ or $13k + 6$

Table 1: Independence numbers of the strong product of three moderate sized odd cycles

The results presented in Table 1 were obtained mostly using backtracking algorithms. For instance, the computations for $\alpha(C_7 \boxtimes C_7 \boxtimes C_9)$ took about one week. In [25] it was also conjectured that the independence numbers of $C_7 \boxtimes C_7 \boxtimes C_{2k+1}$ and $C_7 \boxtimes C_9 \boxtimes C_{2k+1}$ reach the upper bound from Table 1 if k is large enough. This is indeed true for the first case, as follows from the main result of this section:

Theorem 6 *Let $k \geq 3$. Then:*

$$\alpha(C_7 \boxtimes C_7 \boxtimes C_{2k+1}) = \begin{cases} 33; & k = 3, \\ 10k + 4; & 4 \leq k \leq 9, \\ 10k + 5; & k \geq 10. \end{cases}$$

In the rest of the section we prove Theorem 6. It is well-known that for any graph G and $n \in \mathbb{N}$, $\alpha(G \boxtimes K_n) = \alpha(G)$. Hence:

Corollary 7 $\alpha(C_7 \boxtimes C_7 \boxtimes K_2) = \alpha(C_7 \boxtimes C_7) = 10$.

We will also need the following lemma from [24]:

Lemma 8 *For any graph G and any $k \in \mathbb{N}$,*

$$\alpha(C_{2k+3} \boxtimes G) \geq \lceil \frac{2k+3}{2k+1} \alpha(C_{2k+1} \boxtimes G) \rceil.$$

Since $G = C_7 \boxtimes C_7 \boxtimes C_{2k+1} = (C_7 \boxtimes C_7) \boxtimes C_{2k+1}$, we can consider G as the rotagraph $\omega_{2k+1}(C_7 \boxtimes C_7; X)$ where X is defined in the obvious way. We now consider those functions f of $\mathcal{I}_2(\omega_{2k+1}(C_7 \boxtimes C_7; X))$ for which $|f^{-1}(1)| = 10k + 5$. From Corollary 7 we infer that $|f_i^{-1}(1)| = 5$. By Corollary 2, $\omega_{2k+1}(C_7 \boxtimes C_7; X)$ has an independent set with $10k + 5$ independent vertices if and only if $D = D_1(C_7 \boxtimes C_7; X)$ contains a directed closed walk of length $2k + 1$.

We have first computed the directed graph D , giving us $980 \cdot \binom{10}{5} = 246960$ vertices of D . (Recall from [25] that $C_7 \boxtimes C_7$ has 980 independent sets with 10 vertices if its vertices have fixed labels.) Using depth-first search we then discovered that D contains an odd cycle. In order to establish the smallest k , such that $\alpha(C_7 \boxtimes C_7 \boxtimes C_{2k+1}) = 10k + 5$, a shortest odd cycle was to be found. We used a construction described by Grötschel, Lovász and Schrijver in [10]. An auxiliary bipartite graph $\Gamma = ((V^1, V^2), E)$ was constructed as follows. Each vertex $u \in D$ was split into two vertices u^1 and u^2 , with u^i included in V^i ($i = 1, 2$). For each arc uv of D we have added the arcs u^1v^2 and u^2v^1 to E . Note that a shortest path from u^1 to u^2 in Γ corresponds to a shortest cycle of odd length in D containing u . Thus we have constructed Γ and determined the length of a shortest path from u^1 to u^2 for every $u \in D$. The program found out that a shortest cycle in D is of length 21. The corresponding set with 105 independent vertices is depicted in Figure 6.




Figure 6: 105 independent vertices of $C_7 \boxtimes C_7 \boxtimes C_{21}$

In this way we have established the theorem for $k = 10$. The result for $k > 10$ follows from Lemma 8.

6 Concluding remarks

In this paper we have proposed a dynamic algorithm approach for computing graph properties of rotagraphs. Essentially the same approach can be used for computing graph properties of fasciagraphs. For instance, in the case of 1-local or 2-local properties, the only difference being that instead of searching for directed closed walks of length n in $D_d(G; X)$ we need to look for directed walks of length n .

Suppose we consider a d -local, hereditary property \mathcal{P}_q with $d > 2$. Then one might select the fasciagraphs $\Psi_{d+1}(G; X)$ as the basic building stone of the rotagraphs $\omega_n(G; X)$ and combine functions restricted to these fasciagraphs to search for functions on rotagraphs. In this way a corresponding directed graph D can be introduced such that $\mathcal{P}_q(\omega_n(G; X))$ is nonempty if and only if D contains a directed closed walk of length n , a result analogous to Corollary 2. Of course, it depends on the functions considered which approach would be more efficient. In particular, if the approach via fasciagraphs $\Psi_d(G; X)$ would be used, the number of the corresponding functions on $\Psi_d(G; X)$ must be of a moderate size.

Not all interesting graph functions are d -local and hereditary. For instance, we have observed that the functions from \mathcal{D}_2 describing dominating sets are not hereditary. Nevertheless, a slightly altered approach can still be used to obtain new exact domination numbers of the Cartesian product of two cycles. It will be described in a subsequent paper [26].

Finally, we wish to add that in the case of rotagraphs, the concept of a d -local property seems to be most naturally defined in terms of the rotagraph's fibers. However, in a more general situation where we have no fiber structure, it might have been natural to define a graph property \mathcal{P}_q to be d -local, if for any graph (from a certain class of graphs) G , a function $f \in \mathcal{F}_q(G)$ belongs to $\mathcal{P}_q(G)$ if and only if for any vertex u of G the function f , restricted to disk $N_d[u]$ of radius d around u , belongs to $\mathcal{P}_q(\langle N_d[u] \rangle)$.

References

- [1] D. Babić, A. Graovac, B. Mohar and T. Pisanski, The matching polynomial of a polygraph, *Discrete Appl. Math.* 15 (1986) 11–24.
- [2] L.D. Baumert, R.J. McEliece, E. Rodemich, H.C. Rumsey, R. Stanley, H. Taylor, A combinatorial packing problem, in: *Computers in Algebra and Number Theory* (Proc. SIAM-AMS Sympos. Appl. Math.), 1971, Amer. Math. Soc., Providence, R.I., 97–108.
- [3] A.A. Bertossi, M.C. Pinotti, R.B. Tan, Channel assignment with separation for interference avoidance in wireless networks, preprint.
- [4] G. J. Chang and D. Kuo, The (2,1)-labeling problem on graphs, *SIAM J. Discrete Math.* 9 (1996) 309–316.

- [5] J. Georges and D. W. Mauro, Generalized vertex labelings with a condition at distance two, *Congr. Numer.* 109 (1995) 141–159.
- [6] J. Georges and D. W. Mauro, On the size of graphs labeled with a condition at distance two, *J. Graph Theory* 22 (1996) 47–57.
- [7] J. Georges and D. W. Mauro, Some results of λ_k^j -numbers of the products of complete graphs, *Congr. Numer.* 140 (1999) 141–160.
- [8] J. P. Georges, D. W. Mauro and M. I. Stein, Labeling products of complete graphs with a condition at distance two, *SIAM J. Discrete Math.* 14 (2001) 28–35.
- [9] J. Georges, D.W. Mauro and M. Whittlesey, Relating path covering to vertex labelings with a condition at distance two, *Discrete Math.* 135 (1994) 103–111.
- [10] M. Grötschel, L. Lovász and A. Schrijver. *Geometric Algorithms and Combinatorial Optimization.* Springer-Verlag, Berlin, 1988.
- [11] C.D. Godsil, Problems in algebraic combinatorics, *Electron. J. Combin.* 2 (1995) 20 pp.
- [12] J.R. Griggs and R.K. Yeh, Labelling graphs with a condition at distance two, *SIAM J. Discrete Math.* 5 (1992) 586–595.
- [13] R.S. Hales, Numerical invariants and the strong product of graphs, *J. Combin. Theory Ser. B* 15 (1973) 146–155.
- [14] J. Van den Heuvel, R. A. Leese and M. A. Shepherd, Graph labeling and radio channel assignment, *J. Graph Theory* 29 (1998) 263–283.
- [15] W. Imrich, S. Klavžar, *Product Graphs: Structure and Recognition*, John Wiley & Sons, New York, 2000.
- [16] P.K. Jha, Optimal $L(2,1)$ -labeling of Cartesian products of cycles, with an application to independent domination, *IEEE Trans. Circ. Syst.-I: Fund. Theory and Appl.* 47 (2000) 1531–1534.
- [17] P.K. Jha, Optimal $L(2,1)$ -labeling of strong products of cycles, *IEEE Trans. Circ. Syst.-I: Fund. Theory and Appl.* 48 (2001) 498–500.
- [18] P.K. Jha, A. Narayanan, P. Sood, V. Sunder, On $L(2,1)$ -labeling of the Cartesian product of a cycle and a path, *Ars Combin.* 55 (2000) 81–89.
- [19] M. Juvan, B. Mohar and J. Žerovnik, Distance-related invariants on polygraphs, *Discrete Appl. Math.* 80 (1997) 57–71.
- [20] S. Klavžar and J. Žerovnik, Algebraic approach to fasciagraphs and rotagraphs, *Discrete Appl. Math.* 68 (1996) 93–100.

- [21] D. Liu and R. K. Yeh, On distance two labelings of graphs, *Ars Combin.* 47 (1997) 13–22.
- [22] D. Sakai, Labeling chordal graphs: Distance two condition, *SIAM J. Discrete Math.* 7 (1994) 133–140.
- [23] C.E.Shannon, The zero-error capacity of a noisy channel, *IRE Trans. Inform. Theory* 2 (1956) 8–19.
- [24] E. Sonnemann and O. Krafft, Independence numbers of product graphs, *J. Combin. Theory Ser. B* 17 (1974) 133–142.
- [25] A.Vesel, The independence number of the strong product of cycles, *Comput. Math. Appl.* 36 (1998) 9–21.
- [26] A.Vesel, On the domination number of the Cartesian product of two cycles, in preparation.
- [27] A.Vesel and J. Žerovnik, Improved lower bound on the Shannon capacity of C_7 , *Inform. Proc. Lett.*, to appear.
- [28] M. A. Weiss, *Data Structures and Algorithm Analysis*, The Benjamin/Cummings Publishing Company, Redwood City, 1995.
- [29] M. A. Whittlesey, J. P. Georges, and D. W. Mauro, On the λ -number of Q_n and related graphs, *SIAM J. Discrete Math.* 8 (1995) 499–506.
- [30] J. Žerovnik, Deriving formulas for domination numbers of fasciagraphs and rotagraphs, in: *Fundamentals of Computation Theory, 12th International Symposium FCT'99*, Iasi, Romania (G. Ciobanu, Gh. Paun, eds.) *Lecture Notes in Computer Science* 1684, Springer-Verlag, 1999, pp. 559–568.