

UNIVERSITY OF LJUBLJANA  
INSTITUTE OF MATHEMATICS, PHYSICS AND MECHANICS  
DEPARTMENT OF MATHEMATICS  
JADRANSKA 19, 1000 LJUBLJANA, SLOVENIA

**Preprint series, Vol. 39 (2001), 788**

ON A LIST-COLORING  
PROBLEM

Sylvain Gravier      Frédéric Maffray  
Bojan Mohar

ISSN 1318-4865

November 23, 2001

Ljubljana, November 23, 2001

# On a list-coloring problem\*

Sylvain Gravier<sup>†</sup>      Frédéric Maffray<sup>†</sup>      Bojan Mohar<sup>‡</sup>

November 8, 2001

## Abstract

We study the function  $f(G)$  defined for a graph  $G$  as the smallest integer  $k$  such that the join of  $G$  with a stable set of size  $k$  is not  $|V(G)|$ -choosable. This function was introduced recently in order to describe extremal graphs for a list-coloring version of a famous inequality due to Nordhaus and Gaddum [1]. Some bounds and some exact values for  $f(G)$  are determined.

## 1 Introduction

We consider undirected, finite, simple graphs. A *coloring* of a graph  $G = (V, E)$  is a mapping  $c : V \rightarrow \{1, 2, \dots\}$  such that  $c(u) \neq c(v)$  for every edge  $uv \in E$ . If  $|c(V)| \leq k$ , then  $c$  is also said to be a  $k$ -*coloring*. The *chromatic number*  $\chi(G)$  is the smallest integer  $k$  such that  $G$  admits a  $k$ -coloring. A graph is  $k$ -*colorable* if it admits a  $k$ -coloring.

Vizing [4], as well as Erdős, Rubin and Taylor [2] introduced a variant of the coloring problem as follows. Suppose that each vertex  $v$  is assigned a list  $L(v) \subseteq \{1, 2, \dots\}$  of allowed colors; we then want to find a coloring  $c$  such that  $c(v) \in L(v)$  for all  $v \in V$ . If such a coloring exists, we say that  $G$  is  $L$ -*colorable* and that  $c$  is an  $L$ -*coloring* of  $G$ . The graph is  $k$ -*choosable* if  $G$  is  $L$ -colorable for every assignment  $L$  that satisfies  $|L(v)| \geq k$  for all  $v \in V$ . The *choice number* or *list-chromatic number*  $Ch(G)$  of  $G$  is the smallest  $k$  such that  $G$  is  $k$ -choosable. Clearly, every graph satisfies  $Ch(G) \geq \chi(G)$ .

Let  $G_1, G_2$  be two vertex-disjoint graphs. The graph  $G_1 + G_2 = (V(G_1) \cup V(G_2), E(G_1) \cup E(G_2) \cup \{xy \mid x \in V(G_1), y \in V(G_2)\})$  is called the *join* of  $G_1$  and  $G_2$ . It is easy to see that  $\chi(G_1 + G_2) = \chi(G_1) + \chi(G_2)$  for

---

\*This research was partially supported by the Proteus Project 00874RL

<sup>†</sup>CNRS, Laboratoire Leibniz, 46 avenue Félix Viallet, 38031 Grenoble Cédex, France

<sup>‡</sup>Dept. of Mathematics, University of Ljubljana, Slovenia

any two vertex-disjoint graphs  $G_1, G_2$ . So, the chromatic number has a straightforward behavior with respect to the join operation. On the other hand, the choice number does not behave so simply. For instance, if  $G_1$  and  $G_2$  are edgeless graphs on  $n$  and  $n^n$  vertices, respectively, then obviously  $Ch(G_1) = Ch(G_2) = 1$ , but it is easy to check (see [3]) that  $Ch(G_1 + G_2) = n + 1$ , i.e., the complete bipartite graph  $K_{n, n^n}$  is not  $n$ -choosable.

Let us denote by  $S_k$  the edgeless graph on  $k$  vertices. Since the complete bipartite graph  $K_{n, n^n}$  is not  $n$ -choosable, if  $H$  is any graph on  $n$  vertices then  $Ch(H + S_{n^n}) > n$ . We can therefore define  $f(H)$  as the smallest integer  $k$  such that  $Ch(H + S_k) > |V(H)|$ . The fact from [3] that  $K_{n, n^n}$  is not  $n$ -choosable and is minimal with that property means that  $f(S_n) = n^n$ . It is easy to see that  $f(K) = 1$  for every complete graph  $K$ . Obviously, if  $e \in E(G)$ , then  $f(G - e) \geq f(G)$ . This implies:

$$\text{If } G \text{ is any graph on } n \text{ vertices, then } 1 \leq f(G) \leq n^n. \quad (1)$$

The definition of  $f(G)$  was motivated by the determination of extremal graphs for the inequality  $Ch(G) + Ch(\overline{G}) \leq |V(G)| + 1$  (see [1]). Here we would like to examine in more detail the problem of evaluating and computing  $f(G)$ .

An alternative definition for  $f(G)$  can be given as follows. Let  $G = (V, E)$  be a graph on  $n$  vertices, and let  $\mathcal{L}(G)$  be the set of assignments  $L : V \rightarrow \mathcal{P}(\{1, 2, \dots\})$  that satisfy:

- (i)  $|L(v)| \geq n, \forall v \in V$ , and
- (ii)  $L(u) \cap L(v) = \emptyset$  if  $u, v \in V, uv \notin E$ .

Clearly, for every  $L \in \mathcal{L}(G)$ , there exists at least one  $L$ -coloring of  $G$ , because of (i). Moreover, by (ii) every  $L$ -coloring  $c$  of  $G$  uses exactly  $n$  colors; we denote by  $c(V)$  the set of  $n$  colors used by  $c$ . We now write:

$$\mathcal{C}(L) = \{c(V) \mid c \text{ is an } L\text{-coloring of } G\}. \quad (2)$$

Now define  $f'(G) = \min\{|\mathcal{C}(L)| : L \in \mathcal{L}(G)\}$ .

**Lemma 1** *For every graph  $G$ , we have  $f(G) = f'(G)$ .*

**Proof.** Assume  $G$  has  $n$  vertices, and write  $f(G) = k$ . By the definition of  $f(G)$ , we have  $Ch(G + S_k) \geq n + 1$ . Thus there exists a list assignment

$L$  on  $V(G + S_k)$  with  $|L(v)| \geq n$  ( $\forall v \in V(G + S_k)$ ) and such that  $G + S_k$  is not  $L$ -colorable. If there were non-adjacent vertices  $u, v \in V(G)$  such that  $L(u) \cap L(v) \neq \emptyset$ , then we could assign a color from  $L(u) \cap L(v)$  to  $u$  and  $v$  and then extend this greedily to an  $L$ -coloring of  $G$  and then to an  $L$ -coloring of  $G + S_k$ , a contradiction. It follows that the restriction of  $L$  to  $G$  satisfies (i) and (ii). Furthermore, whenever  $c$  is an  $L$ -coloring of  $G$ , the set  $c(V(G))$  must appear as  $L(s)$  for at least one  $s \in S_k$ , for otherwise this  $L$ -coloring  $c$  of  $G$  could obviously be extended to an  $L$ -coloring of  $G + S_k$ , a contradiction. Hence  $|\mathcal{C}(L)| \leq k$ . The definition of  $f'$  implies  $f'(G) \leq k$ , i.e.,  $f'(G) \leq f(G)$ .

Conversely, assume that  $L$  is a list assignment on  $G$  such that  $L \in \mathcal{L}(G)$  and  $|\mathcal{C}(L)| = f'(G) = j$ . Write  $\mathcal{C}(L) = \{C_1, \dots, C_j\}$  and let  $S_j = \{s_1, \dots, s_j\}$  be a stable set of size  $j$ . Let  $L'$  the list assignment defined by  $L'(v) = L(v)$  for all  $v \in V(G)$  and  $L'(s_i) = C_i$  ( $i = 1, \dots, j$ ). Observe that, by (ii),  $|L'(u)| \geq n$  for all  $u \in V(G + S_j)$ . Clearly  $G + S_j$  is not  $L'$ -colorable, so  $f(G) \leq j$ , i.e.,  $f(G) \leq f'(G)$ .  $\square$

Using Lemma 1, it is possible to compute  $f(G)$  for some small graphs, but in general the computation is difficult even for graphs with a simple structure. For example, one can establish that  $f(C_4) = 36$ , but we need a tedious case analysis to show that  $f(C_5) = 500$ .

**Theorem 1** *If  $G$  has  $n$  vertices and is not a complete graph, then  $f(G) \geq n^2$ .*

**Proof.** We will prove, by induction on  $n$ , that if  $u, v$  are non-adjacent vertices of  $G$  and  $L \in \mathcal{L}(G)$ , then  $f'(G) \geq |L(u)||L(v)|$ . This statement clearly implies the theorem. For  $n = 2$ , the statement is obvious. Now, assume that  $n \geq 3$ , and write  $n_1 = |L(u)|$  and  $n_2 = |L(v)|$ . Pick any  $z \in V \setminus \{u, v\}$  and pick any color, say 1, in  $L(z)$ . We may assume by (ii) that  $1 \notin L(v)$ . Define:

$$\mathcal{C}_1(L) = \{c(V) \mid c \text{ is an } L\text{-coloring of } G \text{ with } c(z) = 1\},$$

$$\overline{\mathcal{C}}_1(L) = \{c(V) \mid c \text{ is an } L\text{-coloring of } G \text{ with } 1 \notin c(V)\}.$$

Clearly,  $\mathcal{C}(L) \supseteq \mathcal{C}_1(L) \cup \overline{\mathcal{C}}_1(L)$  and  $\mathcal{C}_1(L) \cap \overline{\mathcal{C}}_1(L) = \emptyset$ . Thus  $|\mathcal{C}(L)| \geq |\mathcal{C}_1(L)| + |\overline{\mathcal{C}}_1(L)|$ . Let us now evaluate these numbers.

On one hand, we have  $|\mathcal{C}_1(L)| \geq (n_1 - 1)n_2$  by the induction hypothesis applied to the graph  $G - z$  with the list assignment  $L_1 \in \mathcal{L}(G - z)$  determined by  $L_1(w) = L(w) \setminus \{1\}$  for each  $w \in V(G - z)$ .

On the other hand, we claim that  $|\overline{\mathcal{C}}_1(L)| \geq n_2$ . Indeed, fix an  $L$ -coloring  $\gamma$  of the subgraph  $G \setminus \{u, v\}$  that does not use color 1. Such a coloring exists because that subgraph has  $n - 2$  vertices while  $L_1$  assigns lists of size at least  $n - 1$  by (i). Write  $t_1 = |L(u) \cap \gamma(V \setminus \{u, v\})|$  and  $t_2 = |L(v) \cap \gamma(V \setminus \{u, v\})|$ . Write  $\lambda_1 = n_1 - (t_1 + 1)$  and  $\lambda_2 = n_2 - t_2$ . Since color 1 is not in  $L(v)$  (but possibly is in  $L(u)$ ),  $\gamma$  can be extended to an  $L$ -coloring of  $G$  in at least  $\lambda_1 \lambda_2$  ways, and each of these uses a different set of colors  $\gamma(V) \in \overline{\mathcal{C}}_1(L)$ . Since  $\lambda_1 > 0$ ,  $\lambda_2 > 0$ , and  $\lambda_1 + \lambda_2 \geq n_2 + 1$ , we have  $|\overline{\mathcal{C}}_1(L)| \geq \lambda_1 \lambda_2 \geq n_2$ .

Now,  $|\mathcal{C}_1(L)| \geq (n_1 - 1)n_2$  and  $|\overline{\mathcal{C}}_1(L)| \geq n_2$  imply  $|\mathcal{C}(L)| \geq n_1 n_2$ .  $\square$

We observe that the bound given in the preceding theorem is tight, i.e., for any  $n \geq 2$ , there exists a graph  $G$  on  $n$  vertices with  $f(G) = n^2$ . Indeed, consider the graph  $K_n - E(K_{1,i})$  obtained from a complete graph on  $n$  vertices by removing  $i$  edges incident to one given vertex  $u$  ( $1 \leq i \leq n - 1$ ):

**Claim 1**  $f(K_n - E(K_{1,i})) = n^2$ .

**Proof.** By Theorem 1, we have  $f(K_n - E(K_{1,i})) \geq n^2$ , so we need only to prove that  $f(K_n - E(K_{1,i})) \leq n^2$ . For this purpose, assign to the vertex  $u$  the list  $\{1, 2, \dots, n\}$  and to all other vertices of the graph the list  $\{n + 1, \dots, 2n\}$ . This yields a list assignment  $L \in \mathcal{L}(G)$ . It is easy to check that  $|\mathcal{C}(L)| = n^2$ , hence  $f(G) \leq n^2$ .  $\square$

We do not know of any graph  $G$  other than  $K_n - E(K_{1,i})$  that satisfies  $f(G) = |V(G)|^2$ .

## 2 The significance of clique partitions

Given a graph  $G = (V, E)$ , a *clique partition* of  $G$  is a set  $Q = \{Q_1, \dots, Q_p\}$  of pairwise disjoint, non-empty cliques such that  $V = Q_1 \cup \dots \cup Q_p$ . Let  $n = |V|$  and  $q_i = |Q_i|$ ,  $i = 1, \dots, p$ . Then we write

$$w(Q) = \prod_{i=1}^p \binom{n}{q_i}$$

and

$$w(G) = \min\{w(Q) \mid Q \text{ is a clique partition of } G\}.$$

**Theorem 2** *For every graph  $G$ , we have  $f(G) \leq w(G)$ .*

**Proof.** Write  $n = |V|$ . Consider a clique partition  $Q = \{Q_1, \dots, Q_p\}$  of  $G$ , and make a list assignment  $L$  as follows: to each vertex of  $Q_i$  assign a list  $L_i$  of  $n$  colors, so that  $L_i \cap L_j = \emptyset$  whenever  $1 \leq i < j \leq p$ . Clearly,  $L \in \mathcal{L}(G)$ . Moreover, any  $L$ -coloring of  $G$  consists in assigning  $|Q_1|$  colors from  $L_1$  to the vertices of  $Q_1$ ,  $|Q_2|$  colors from  $L_2$  to the vertices of  $Q_2$ , etc. It follows that  $|\mathcal{C}(L)| = w(Q)$ . Therefore,  $f'(G) \leq w(Q)$ . Since  $Q$  is an arbitrary clique partition, Lemma 1 implies that  $f(G) = f'(G) \leq w(G)$ .  $\square$

**Claim 2** *If  $G$  is a disjoint union of cliques, then  $f(G) = w(G)$ .*

**Proof.** By the preceding theorem, we need only prove  $f(G) \geq w(G)$ . Assume  $G$  is the union of cliques  $Q_1, \dots, Q_p$ . Consider any list assignment  $L \in \mathcal{L}(G)$ . Let us denote by  $L^i$  the restriction of  $L$  to the subgraph of  $G$  induced by  $Q_i$  ( $i = 1, \dots, p$ ). Note that the colors assigned by  $L^i$  to any vertex in  $Q_i$  are different from the colors assigned by  $L^j$  to any vertex in  $Q_j$  whenever  $i \neq j$ , by (ii). Thus  $|\mathcal{C}(L)| = |\mathcal{C}(L^1)| \cdots |\mathcal{C}(L^p)|$ . Every  $L^i$ -coloring of  $Q_i$  can be obtained by choosing among at least  $n$  colors for the first vertex of  $Q_i$ , then among at least  $n - 1$  available colors for the second vertex, etc. This way, a given set of  $|Q_i|$  colors used in such a coloring occurs at most  $|Q_i|!$  times. Thus,

$$|\mathcal{C}(L^i)| \geq \frac{n(n-1) \cdots (n - |Q_i| + 1)}{|Q_i|!} = \binom{n}{|Q_i|}.$$

Consequently,  $|\mathcal{C}(L)| \geq w(Q) \geq w(G)$ . Since  $L$  was an arbitrary element of  $\mathcal{L}(G)$ , the result follows.  $\square$

The preceding fact shows that the inequality in Theorem 2 is best possible and motivates the following conjecture.

**Conjecture 1** *For every graph  $G$ , we have  $f(G) = w(G)$ .*

We note that if  $G$  is a triangle-free graph on  $n$  vertices, a clique partition  $Q$  consists of some cliques of size two (which form a matching) and some cliques of size one. If  $p_2$  is the number of cliques of size two, we see that  $w(Q) = \binom{n}{2}^{p_2} n^{n-2p_2}$ ; this number is minimized when  $p_2$  is maximized, i.e., when the cliques of size two in  $Q$  form a matching of  $G$  of maximum size. We denote by  $\mu(G)$  the size of a maximum matching. This leads us to:

**Conjecture 2** *For every triangle-free graph  $G$ ,  $f(G) = \binom{n}{2}^{\mu(G)} n^{n-2\mu(G)}$ .*

This conjecture suggests that the computation of  $f(G)$  should be tractable for triangle-free graphs. We have not been able to prove this second conjecture, not even in the case of trees. The following lemma will help us settle a special case.

For a graph  $G = (V, E)$  and two adjacent vertices  $u, v$  of  $G$ , define  $\mathcal{L}_{uv}(G) = \{L \in \mathcal{L}(G) \mid L(u) = L(v)\}$ .

**Lemma 2** *Let  $G$  be a graph and  $uv$  an edge of  $G$  such that  $u$  is of degree 1 and  $v$  is of degree at most 2 in  $G$ . Then, for each  $L \in \mathcal{L}(G)$ , there exists  $L' \in \mathcal{L}_{uv}(G)$  such that  $L'(x) = L(x)$ , for every  $x \in V \setminus \{u, v\}$  and  $|\mathcal{C}(L')| \leq |\mathcal{C}(L)|$ .*

**Proof.** Write  $U = \bigcup\{L(x) \mid x \in V \setminus \{u, v\}\}$  and observe that  $L(u)$  is disjoint from  $U$ . If  $L(v)$  too is disjoint from  $U$ , we set  $L'(u) = L'(v) = L(u)$ , and we set  $L'(x) = L(x)$  for  $x \in V \setminus \{u, v\}$ . Then it is easy to check that  $|\mathcal{C}(L')| \leq |\mathcal{C}(L)|$ .

Now assume that  $L(v)$  is not disjoint from  $U$ . Since  $L$  satisfies (ii), this means that  $v$  has another neighbour  $w$ , and that  $L(v) \cap U = L(v) \cap L(w)$ . Write  $B = L(u) \cap L(v)$  and  $C = L(v) \cap L(w)$ , and then  $A = L(u) \setminus B$ ,  $P = L(v) \setminus (B \cup C)$ , and  $D = L(w) \setminus C$ . Thus we have  $L(u) = A \cup B$ ,  $L(v) = B \cup C \cup P$ ,  $L(w) = C \cup D$ , with  $A \cap B = B \cap C = B \cap P = C \cap P = C \cap D = \emptyset$ , and  $C \neq \emptyset$ .

We can assume that  $|A| \leq |C \cup P|$ . Indeed, if  $|A| > |C \cup P|$ , we replace  $L$  by the assignment  $L^*$  obtained by removing  $|A| - |C \cup P|$  elements of  $A$  from  $L(u)$  and by setting  $L^*(x) = L(x)$  for  $x \in V \setminus \{u\}$ . Clearly,  $|\mathcal{C}(L^*)| \leq |\mathcal{C}(L)|$ . The corresponding sets  $A^*, C^*, P^*$  of  $L^*$  satisfy  $|A^*| = |C^* \cup P^*|$  so we can work with  $L^*$  instead of  $L$ .

We fix a one-to-one mapping  $a \mapsto \bar{a}$  from  $A$  to  $C \cup P$ .

Define  $L'$  by  $L'(u) = L'(v) = L(u) = A \cup B$  and  $L'(x) = L(x)$  if  $x \in V \setminus \{u, v\}$ . We claim that  $L'$  satisfies the conclusion of the lemma. Clearly,  $L' \in \mathcal{L}_{uv}(G)$ .

Let  $\gamma'$  be an  $L'$ -coloring of  $G$ . We denote elements of  $A$  and  $B$  by the corresponding lowercase letters, and we write, e.g.,  $\gamma'(u, v) = (a, b)$  as a shorthand for  $\gamma'(u) = a \in A$ ,  $\gamma'(v) = b \in B$ . Observe that for  $\gamma'(u, v)$ , there are four possibilities:  $(a_1, a_2)$ ,  $(a, b)$ ,  $(b, a)$ , and  $(b_1, b_2)$ . Define a mapping  $\gamma$  by  $\gamma(x) = \gamma'(x)$  for all  $x \in V \setminus \{u, v\}$ . We extend  $\gamma$  to an  $L$ -coloring of  $G$  as follows:

If  $\gamma'(u, v)$  is either  $(a, b)$  or  $(b, a)$ , set  $\gamma(u, v) = (a, b)$ .

If  $\gamma'(u, v) = (b_1, b_2)$ , set  $\gamma(u, v) = (b_1, b_2)$ .

If  $\gamma'(u, v) = (a_1, a_2)$ , set  $\gamma(u, v) = (a_1, \bar{a}_2)$  if  $\bar{a}_2 \neq \gamma'(w)$ ; otherwise set  $\gamma(u, v) = (a_2, \bar{a}_1)$ .

Clearly,  $\gamma$  is an  $L$ -coloring. Moreover, it is a routine matter to check that whenever  $\gamma', \delta'$  are two  $L'$ -colorings with  $\gamma'(V) \neq \delta'(V)$  then the corresponding  $L$ -colorings  $\gamma, \delta$  satisfy  $\gamma(V) \neq \delta(V)$ . This implies that  $|\mathcal{C}(L')| \leq |\mathcal{C}(L)|$ .  $\square$

As an application, consider the class  $\mathcal{B}$  of trees obtained from the trees on one or two vertices by repeating the following construction: add a vertex  $v$  of degree one, and then add a vertex  $u$  adjacent only to  $v$ .

**Corollary 1** *If  $G$  is an  $n$ -vertex graph in  $\mathcal{B}$ , then  $f(G) = \binom{n}{2}^{\mu(G)} n^{n-2\mu(G)}$ .*

**Proof.** Let  $v_1, u_1, \dots, v_k, u_k$  be the vertices used in the recursive construction of  $G$ . Note that  $u_k$  is pendant in  $G$ , hence  $v_k u_k$  belongs to a maximum matching of  $G$ . Recursively this implies that  $M = \{v_1 u_1, \dots, v_k u_k\}$  is a maximum matching of  $G$ , hence  $k = \mu(G)$ . Consider any  $L \in \mathcal{L}(G)$ . Applying the preceding lemma repeatedly, we obtain an assignment  $L' \in \mathcal{L}(G)$  which satisfies  $|\mathcal{C}(L')| = \binom{n}{2}^k n^{n-2k} \leq |\mathcal{C}(L)|$ .  $\square$

## References

- [1] S. Dantas, S. Gravier, F. Maffray, Extremal graphs for the list-coloring version of a theorem of Nordhaus and Gaddum, Research Report 18, Laboratoire Leibniz-IMAG, Grenoble, 2000.
- [2] P. Erdős, A. L. Rubin, H. Taylor, Choosability in graphs, *Congr. Numer.* 26 (1979), 125–157.
- [3] N. V. R. Mahadev, F. S. Roberts, P. Santhanakrishnan, 3-choosable complete bipartite graphs, RUTCOR Research Report 49-91, Rutgers Univ., NJ, USA (1991).
- [4] V. G. Vizing, Vertex colourings with given colors (in Russian), *Metody Discret. Analiz.* 29 (1976), 3–10.