

UNIVERSITY OF LJUBLJANA  
INSTITUTE OF MATHEMATICS, PHYSICS AND MECHANICS  
DEPARTMENT OF MATHEMATICS  
JADRANSKA 19, 1000 LJUBLJANA, SLOVENIA

**Preprint series, Vol. 40 (2002), 824**

ON CUBIC AND  
EDGE-CRITICAL ISOMETRIC  
SUBGRAPHS OF HYPERCUBES

C. Paul Bonnington      Sandi Klavžar  
                                         Alenka Lipovec

ISSN 1318-4865

May 6, 2002

Ljubljana, May 6, 2002

# On cubic and edge-critical isometric subgraphs of hypercubes

C. Paul Bonnington\*  
Department of Mathematics  
University of Auckland  
Auckland, New Zealand  
p.bonnington@auckland.ac.nz

Sandi Klavžar†  
Department of Mathematics, University of Maribor  
Koroška cesta 160, 2000 Maribor, Slovenia  
sandi.klavzar@uni-lj.si

Alenka Lipovec  
Department of Education, University of Maribor  
Koroška cesta 160, 2000 Maribor, Slovenia  
alenka.lipovec@uni-mb.si

## Abstract

All cubic partial cubes (i.e., cubic isometric subgraphs of hypercubes) up to 30 vertices and all edge-critical partial cubes up to 14 vertices are presented. The lists of graphs were confirmed by computer search to be complete. Non-trivial cubic partial cubes on 36, 42, and 48 vertices are also constructed.

## 1 Introduction

*Partial cubes* are, by definition, graphs that admit isometric embeddings into hypercubes. They were introduced by Graham and Pollak [10] and first characterized by Djoković [7]. Several additional characterizations followed in [2, 5, 19, 20]. Partial cubes found different applications (see, for example, [6, 8, 13]), while recognition algorithms for these graphs have been developed in [1, 11]. For an extensive presentation of partial cubes we refer the reader to the book [12].

---

\*Supported by the Marsden Fund (Grant Number UOA-825) administered by the Royal Society of New Zealand.

†Supported by the Ministry of Education, Science and Sport of Slovenia under the grant 0101-504.

For the (probably) most important subclass of partial cubes, median graphs, Mulder [17] proved that hypercubes are the only regular median graphs. In other words, the only regular median graphs are Cartesian products of copies of  $K_2$ . This result has been in [4] extended to the so-called “tree-like” partial cubes. Hence, it is natural to ask which graphs are regular partial cubes. (Regular subgraphs of hypercubes are studied in [18]). Despite the fact that the structure of partial cubes has been well clarified by now, this question seems to be a difficult one.

The Cartesian product of two (regular) partial cubes is a (regular) partial cube. Since even cycles are regular partial cubes, one may wonder whether we get all regular partial cubes as Cartesian products of copies of  $K_2$  and even cycles. In particular, are all cubic partial cubes of the form  $C_{2k} \square K_2$ ,  $k \geq 2$ ? This was believed to be true for quite a while, until two sporadic examples appeared: the generalized Petersen graph  $P(10, 3)$  on 20 vertices, cf. [14], and the graph  $B_1$  (see Fig. 1) on 24 vertices from [9], (see also [12]). Calling the graphs  $C_{2k} \square K_2$ ,  $k \geq 2$ , *trivial cubic partial cubes*, we have verified that besides these two graphs there is only one other nontrivial cubic partial cube on at most 30 vertices. The third example, denoted  $B'_1$  (see Fig. 2), has 30 vertices. It can be obtained from the nontrivial partial cube on 24 vertices by the so-called expansion and was also found by computer search.




Figure 1: Graphs  $B_1$  and  $B_2$

*Edge-critical partial cubes* are partial cubes  $G$  for which  $G - e$  is not a partial cube for all edges  $e$  of  $G$ . The 3-cube and the subdivision graph of  $K_4$  are the only edge-critical partial cubes on at most 10 vertices [15].

In this note we present all cubic partial cubes up to 30 vertices and all edge-critical partial cubes up to 14 vertices. The lists of graphs were confirmed by computer search

to be complete. We also give further larger non-trivial cubic partial cubes on 36, 42, and 48 vertices.

## 2 Cubic partial cubes

A graph  $G$  is called *prime* (with respect to the Cartesian graph product) if  $G = G_1 \square G_2$  implies either  $G_1$  or  $G_2$  is the one-vertex graph  $K_1$ .

The Cartesian product of two regular partial cubes is a regular partial cube. Therefore the problem of characterizing regular partial cubes reduces to prime (with respect to the Cartesian product) partial cubes. For the cubic case, this fact leads to the following observation:

**Proposition 2.1** *Let  $G$  be a cubic partial cube. Then either  $G = C_{2n} \square K_2$  for some  $n \geq 2$  or  $G$  is a prime graph.*

**Proof.** Assume  $G = G_1 \square G_2$ , where  $G_1, G_2 \neq K_1$ . As  $G$  is connected, then so are  $G_1$  and  $G_2$ . Since  $G$  is cubic and the degree of  $(u, v) \in V(G_1 \square G_2)$  is the sum of the degrees of  $u \in G_1$  and  $v \in G_2$ , then one of the factors, say  $G_2$ , contains only vertices of degree one or less. Therefore  $G_2 = K_2$ . Furthermore,  $G_1$  must be 2-regular, and hence a cycle. Moreover, it is an even cycle since partial cubes are bipartite graphs.  $\square$

We now construct the nontrivial cubic partial cubes  $B'_1, B''_1, B'''_1$ , and  $B_2$  on 30, 36, 42, and 48 vertices, respectively. The later graph is shown in Fig. 1, while the others are given in in Fig. 2. These graphs can be constructed by expansions from  $B_1$ , and hence we first introduce the concept of expansion.

Let  $G'$  be a connected graph. A *proper cover* consists of two isometric subgraphs  $G'_1, G'_2$  of  $G'$  such that  $G' = G'_1 \cup G'_2$ ,  $G'_0 = G'_1 \cap G'_2$  is a nonempty subgraph, and there are no edges between  $G'_1 \setminus G'_2$  and  $G'_2 \setminus G'_1$ . (The subgraph  $G'_0$  is called the *intersection* of the cover.) The *expansion* of  $G'$  with respect to  $G'_1, G'_2$  is the graph  $G$  constructed as follows: Let  $G_i$  be an isomorphic copy of  $G'_i$ , for  $i = 1, 2$ , and, for any vertex  $u'$  in  $G'_0$ , let  $u_i$  be the corresponding vertex in  $G_i$ , for  $i = 1, 2$ . Then  $G$  is obtained from the disjoint union  $G_1 \cup G_2$ , where for each  $u'$  in  $G'_0$  the vertices  $u_1$  and  $u_2$  are joined by an edge.

Chepoi [5] proved that a graph is a partial cube if and only if it can be obtained from  $K_1$  by a sequence of expansions. This result was later independently obtained in [8] and is analogous to the convex expansion theorem for median graphs [16].

An expansion is called *peripheral* if at least one of the graphs  $G'_1$  or  $G'_2$  is equal to  $G$ . In this situation the other graph equals the intersection, and is necessarily isometric




Figure 2: Graphs  $B_1$ ,  $B'_1$ ,  $B''_1$ , and  $B'''_1$

in  $G$ . We recall from [4] that a regular, prime partial cube on at least three vertices can not be obtained by peripheral expansion from some partial cube.

For the proof of the next result we also need the following concept of isometric dimension. Two edges  $e = xy$  and  $f = uv$  of a graph  $G$  are in the Djoković-Winkler [7, 20] relation  $\Theta$  if  $d_G(x, u) + d_G(y, v) \neq d_G(x, v) + d_G(y, u)$ . Winkler [20] showed that a bipartite graph is a partial cube if and only if  $\Theta = \Theta^*$  (where  $\Theta^*$  denotes the transitive closure of  $\Theta$ ). Thus  $\Theta$  defines an equivalence relation on the edges of a partial cube. The *isometric dimension*,  $\text{idim}(G)$ , of a partial cube  $G$  is defined as the number of its  $\Theta$ -classes.

**Theorem 2.2** *Graphs  $B'_1$ ,  $B''_1$ ,  $B'''_1$ , and  $B_2$  are cubic prime partial cubes.*

**Proof.** We know already that  $B_1$  is a partial cube. Now,  $B'_1, B''_1, B'''_1$ , and  $B_2$  can be obtained from  $B_1, B'_1, B''_1$ , and  $B'''_1$ , respectively, by an expansion. These expansions are schematically explained in Fig. 2 in the following way. A proper cover in each expansion is chosen as follows:  $G'_1$  is induced by the vertices denoted by filled circles,  $G'_2$  is induced by the vertices denoted by filled squares and their intersection is formed by the remaining vertices; that is, the vertices denoted by filled circles surrounded by another circle. It is easy to verify that in this way we really obtain a proper cover; that is,  $G'_1$  and  $G'_2$  are isometric subgraphs of the corresponding graphs  $B_1, B'_1, B''_1$ , and  $B'''_1$ , and there are no edges between  $G'_1 \setminus G'_2$  and  $G'_2 \setminus G'_1$ . Hence, by the theorem of Chepoi the obtained graphs are partial cubes. Clearly, they are cubic.

We now show that these four graphs are prime. Observe first that  $\text{idim}(B_1) = 6$  and therefore  $\text{idim}(B'_1) = 7$ ,  $\text{idim}(B''_1) = 8$ ,  $\text{idim}(B'''_1) = 9$ , and  $\text{idim}(B_2) = 10$ . If any of these four graphs were not prime, then by Proposition 2.1 it would be isomorphic to  $C_{15} \square K_2, C_{18} \square K_2, C_{21} \square K_2$ , and  $C_{24} \square K_2$ , respectively. Two of these graphs are not bipartite, while the isometric dimensions of the other two; that is, of  $C_{18} \square K_2$ , and  $C_{24} \square K_2$ , are 10 and 13.  $\square$

It seems tempting to continue the expansion procedure with  $B_2$  to obtain new cubic partial cubes. However, we were not able to obtain more examples in this way. In particular, the graph that is constructed from  $B_2$  analogously as  $B_2$  is constructed from  $B_1$  is not a partial cube.

### 3 Computer search for cubic and edge-critical partial cubes

Using the Djoković-Winkler relation, we have implemented a recognition algorithm for partial cubes and applied it to all connected bipartite cubic graphs up to 30 vertices. (These graphs were constructed using Brendan McKay's Nauty program [3].) The examination of the entire set of graphs was run concurrently on a cluster of 16 pentium-class machines, and doubled-checked on a 8 processor Sun Sparc server. The obtained results are summarized in the following table:

$n$		$n$		
$< 8$	-	20	$C_{10} \square K_2$	$P(10, 3)$
8	$C_4 \square K_2$	22	-	-
10	-	24	$C_{12} \square K_2$	$B_1$
12	$C_6 \square K_2$	26	-	-
14	-	28	$C_{14} \square K_2$	-
16	$C_8 \square K_2$	30	-	$B'_1$
18	-			

The above table shows that, up to 30 vertices, there are only 3 nontrivial cubic partial cubes.

Recall that a partial cube  $G$  is called edge-critical if for any edge  $e$  of  $G$ ,  $G - e$  is not a partial cube. In [15] it was shown that the 3-cube  $Q_3$  and the subdivision graph of  $K_4$ ,  $S(K_4)$ , are the only edge-critical partial cubes on at most 10 vertices. Moreover, two such graphs on 12 vertices and one on 13 vertices are listed. We have now searched for all edge-critical partial cubes on at most 14 vertices and established the following complete list of edge-critical partial cubes. The computation is a variant of that used for cubic partial cubes; for each connected bipartite graph  $G$  on at most 14 vertices, if  $G$  is determined to be a partial cube, then all of the non-isomorphic graphs obtained by deleting a single edge from  $G$  are tested. Brendan McKay's Nauty program [3] is used to also filter isomorphic graphs from the edge deletions. The results are summarized in the following table.

$n$	$< 8$	8	9	10	11	12	13	14
	-	$Q_3$	-	$S(K_4)$	$E_1$	$E_2, E_3, E_4$	$E_5$	$E_7, \dots, E_{12}$

We note that there is one previously undiscovered graph on each of 11 and 12 vertices ( $E_1$  and  $E_2$  respectively), and six on 14 vertices. These are given in Fig. 3 and 4.




Figure 3: Edge-critical partial cubes on 11, 12, and 13 vertices




Figure 4: Edge-critical partial cubes on 14 vertices

## References

- [1] F. Aurenhammer and J. Hagauer, Recognizing binary Hamming graphs in  $O(n^2 \log n)$  time, *Math. Systems Theory* 28 (1995) 387–395.
- [2] D. Avis, Hypermetric spaces and the Hamming cone, *Canad. J. Math.* 33 (1981) 795–802.
- [3] B. McKay, NAUTY user's guide, Technical report TR-CS-90-02, Comp. Sci. Dept., Australian National University. <http://cs.anu.edu.au/people/bdm/nauty/>
- [4] B. Brešar, W. Imrich, S. Klavžar, Tree-like isometric subgraphs of hypercubes, submitted.

- [5] V. Chepoi,  $d$ -Convexity and isometric subgraphs of Hamming graphs, *Cybernetics* 1 (1988) 6–9.
- [6] V. Chepoi and S. Klavžar, The Wiener index and the Szeged index of benzenoid systems in linear time, *J. Chem. Inform. Comput. Sci.* 37 (1997) 752–755.
- [7] D. Djoković, Distance preserving subgraphs of hypercubes, *J. Combin. Theory Ser. B* 14 (1973) 263–267.
- [8] K. Fukuda and K. Handa, Antipodal graphs and oriented matroids, *Discrete Math.* 111 (1993) 245–256.
- [9] E. Gedeonova, Constructions of  $S$ -lattices, *Order* 7 (1990) 249–266.
- [10] R.L. Graham and H. Pollak, On the addressing problem for loop switching, *Bell System Tech. J.* 50 (1971) 2495–2519.
- [11] W. Imrich, and S. Klavžar, A simple  $O(mn)$  algorithm for recognizing Hamming graphs, *Bull. Inst. Comb. Appl.* 9 (1993) 45–56.
- [12] W. Imrich and S. Klavžar, *Product Graphs: Structure and Recognition*, (Wiley, New York, 2000).
- [13] S. Klavžar and I. Gutman, Wiener number of vertex-weighted graphs and a chemical application, *Discrete Appl. Math.* 80 (1997) 73–81.
- [14] S. Klavžar and A. Lipovec, Partial cubes as subdivision graphs and as generalized Petersen graphs, *Preprint Ser. Dept. Math. Univ. Ljubljana*, vol. 39 (2001) no. 737. Also: *Discrete Math.*, to appear.
- [15] S. Klavžar and A. Lipovec, Edge-critical isometric subgraphs of hypercubes, *Preprint Ser. Dept. Math. Univ. Ljubljana*, vol. 39 (2001) no. 767. Also: *Ars Combin.*, to appear.
- [16] H.M. Mulder, The structure of median graphs, *Discrete Math.* 24 (1978) 197–204.
- [17] H. M. Mulder,  $n$ -Cubes and median graphs, *J. Graph Theory* 4 (1980) 107–110.
- [18] M. Ramras, Regular subgraphs of hypercubes, *Ars Combin.* 52 (1999) 21–32.
- [19] E. Wilkeit, Isometric embeddings in Hamming graphs, *J. Combin. Theory Ser. B* 50 (1990) 179–197.
- [20] P. Winkler, Isometric embeddings in products of complete graphs, *Discrete Appl. Math.* 7 (1984) 221–225.