

UNIVERSITY OF LJUBLJANA
INSTITUTE OF MATHEMATICS, PHYSICS AND MECHANICS
DEPARTMENT OF MATHEMATICS
JADRANSKA 19, 1 111 LJUBLJANA, SLOVENIA

Preprint series, Vol. 43 (2005), 978

TRIANGULATIONS AND THE
HAJÓS CONJECTURE

Bojan Mohar

ISSN 1318-4865

May 20, 2005

Ljubljana, May 20, 2005

Triangulations and the Hajós Conjecture

Bojan Mohar*

Department of Mathematics,
University of Ljubljana,
1000 Ljubljana, Slovenia
`bojan.mohar@fmf.uni-lj.si`

Abstract

The Hajós Conjecture was disproved in 1979 by Catlin. Recently, Thomassen showed that there are many ways that Hajós conjecture can go wrong. On the other hand, he observed that locally planar graphs and triangulations of the projective plane and the torus satisfy Hajós Conjecture, and he conjectured that the same holds for arbitrary triangulations of closed surfaces. In this note we disprove the conjecture and show that also in the case of triangulations, there are many reasons why the Hajós Conjecture can fail.

1 Introduction

Hajós conjecture claims that every graph whose chromatic number is at least k contains a subdivision of K_k , the complete graph of order k . The conjecture has been proved for $k \leq 4$ by Dirac [3], while for $k = 5$, it yields a strengthening of the Four Color Theorem, which is still open. The conjecture was disproved for all $k \geq 7$ by Catlin [2]. Soon after that, Erdős and Fajtlowicz [4] proved that the conjecture is false for almost all graphs.

Recently, Thomassen [11] revived the interest in Hajós conjecture by showing that there is a great variety of reasons why Hajós conjecture can be wrong. At the end of this interesting work, Thomassen observed that the Hajós conjecture could be true to some limited extent. Maybe it holds in the setting under whose influence it was originally formulated (related to the Four Color Conjecture). For instance, it holds for locally planar graphs. Therefore, it seems plausible to propose:

*Supported in part by the Ministry of Education, Science and Sport of Slovenia, Research Project J1-0502-0101-00 and Research Program P1-0297.

Conjecture 1.1 (Thomassen [11]) *Every graph that triangulates some surface satisfies Hajós conjecture.*

Every n -vertex triangulation of a nonplanar surface has at least $3n - 3$ edges. Therefore, it contains a subdivision of K_5 by a theorem of Mader [6]. In particular, possible counterexamples to Conjecture 1.1 must have chromatic number at least 6. Thomassen [11] used known list of 6- and 7-critical graphs on the projective plane and the torus [10] to prove that Conjecture 1.1 holds for triangulations on these two surfaces.

In this note we provide counterexamples to Conjecture 1.1. Additionally, we give some reasons showing that, in certain sense, almost all graphs should be close to some counterexamples.

2 A small counterexample

Let \tilde{H} be the graph with vertex set

$$V = \{v_{i,j} \mid 0 \leq i \leq 4, 0 \leq j \leq 2\} \cup \{w_i \mid 0 \leq i \leq 4\}$$

in which distinct vertices $v_{i,j}$ and $v_{k,l}$ are adjacent if and only if $k \in \{i - 1, i, i + 1\}$ (where $i - 1$ and $i + 1$ are taken modulo 5), and w_i is adjacent to all vertices $v_{i,j}$ and $v_{i+1,j}$, $0 \leq j \leq 2$. In other words, \tilde{H} is composed of 5 copies Q_i ($0 \leq i \leq 4$) of the graph K_7 with vertices $C_i^- = \{v_{i,0}, v_{i,1}, v_{i,2}\}$, $C_i^+ = \{v_{i+1,0}, v_{i+1,1}, v_{i+1,2}\}$, and w_i , and $C_i^+ \subset Q_i$ is identified with $C_{i+1}^- \subset Q_{i+1}$ for $0 \leq i \leq 4$ (indices modulo 5).

The graph \tilde{H} is a counterexample to Conjecture 1.1:

Theorem 2.1 *The graph \tilde{H} triangulates the orientable surface of genus 6 and the nonorientable surface of genus 12. Its chromatic number is 8 and it does not contain a subdivision of K_8 .*




Figure 1: K_7 on the torus

Proof. The graph K_7 triangulates the torus. See Figure 1, where the edge on the left is identified with the rightmost one, and the two cycles of the resulting cylinder are then identified as shown by the labels. Consider 5 such triangulations using Q_0, \dots, Q_4 , such that the triangles C_i^+ and C_i^- of each Q_i are facial; suppose that they correspond to the triangles that are labeled 023 and 156 in Figure 1. By identifying C_i^+ with C_{i+1}^- for $i = 0, 1, 2, 3$, we obtain the connected sum of five tori. Finally, by identifying C_4^+ with C_0^- , the orientable surface of genus 6 is obtained. Clearly, the resulting graph is isomorphic to \tilde{H} and triangulates the surface. Instead of the orientable surface we can get a nonorientable surface of the same Euler characteristic by taking for C_0^- in Q_0 the triangle 165 instead of 156.

Let H be the subgraph of \tilde{H} obtained by removing vertices w_0, \dots, w_4 . Clearly, no three vertices of H are independent. Thus, $\chi(H) \geq \lceil |V(H)|/2 \rceil = 8$. Indeed, it is easy to find an 8-coloring of H , and any such coloring extends to \tilde{H} since the degrees of the removed vertices w_i are equal to 6. Consequently, $\chi(\tilde{H}) = \chi(H) = 8$.

Let us now prove that \tilde{H} does not contain a subdivision of K_8 . Assume, by reductio ad absurdum, that $L \subseteq \tilde{H}$ is such a subgraph. Vertices of degree 7 in L are *branch vertices*; vertices of degree 2 subdivide edges of K_8 . Since $\deg_{\tilde{H}}(w_i) = 6$, no w_i is a branch vertex. Since the neighbors of w_i form a complete subgraph, we may also assume that w_i does not subdivide an edge in L for $i = 0, \dots, 4$. In particular, we conclude that $L \subseteq H$.

Let $B \subseteq V(L)$ be the set of branch vertices and let s be the number of vertices of degree 2 in L . Clearly, $|B| = 8$ and $s \leq |V(H) \setminus B| = 7$. Let $B_i = B \cap C_i^+$ and $b_i = |B_i|$ for $0 \leq i \leq 4$. We may assume that $b_0 = \max_i b_i$ and that, according to this assumption, $b_0 + b_1$ is maximum. Then it is easy to see that $(b_0, b_1) \in \{(3, 3), (3, 2), (3, 1), (2, 2)\}$. We let $b = b_0 + b_1$.

There are $8 - b$ branch vertices that need to be joined to b branch vertices in $B_0 \cup B_1$. Each such vertex $x \in B_2 \cup B_3$ is linked to all vertices in $B_0 \cup B_1$; some of the subdivided edges must contain two or three vertices of degree 2. It is easy to see that for every such x , at least four vertices of degree 2 are needed, with two possible exceptions in the cases when $b_0 = b_1 = 2$ or $b_0 = 3, b_1 = 1$. In any case, we conclude that $s \geq 8$, and this contradiction completes the proof. \square

3 Paley graphs

Let $q = p^r$ be a prime power, $q \equiv 1 \pmod{4}$. The *Paley graph* \mathbf{P}_q is the Cayley graph of the additive group of the finite field $GF(q)$ generated by

all squares, i.e., $V(\mathbf{P}_q) = GF(q)$ and two distinct vertices x, y are adjacent if and only if $x - y = z^2$ for some $z \in GF(q)$. Since $q \equiv 1 \pmod{4}$, $x - y$ is a square if and only if $y - x$ is a square.

Paley graphs have a number of intriguing properties. First of all, they are highly symmetric. Secondly, they are self-complementary. On the other hand, they resemble random graphs a lot. One of their properties which makes them similar to random graphs is the following result of Thomason [9]; see also [1, p. 363].

Theorem 3.1 *Let A be a set of vertices of the Paley graph \mathbf{P}_q . Let $a = |A|$, and let $e(A)$ be the number of edges in the induced subgraph $\mathbf{P}_q(A)$. Then*

$$\left| e(A) - \frac{1}{2} \binom{a}{2} \right| \leq \frac{a(q-a)}{4\sqrt{q}}.$$

This theorem implies that the maximum clique in \mathbf{P}_q has at most \sqrt{q} vertices. Equality is attained if q is an even power of a prime. But if q is a prime, it seems that the maximum cliques in \mathbf{P}_q have far smaller orders. It is believed that their orders are at most polylogarithmic, possibly $o(\log^2 q)$ or even smaller. See [1] for some discussion on this problem.

Let z be a generator of the multiplicative group $GF(q)^*$. It is known that this group is cyclic, so $z^0, z^2, z^4, \dots, z^{q-3}$ are precisely all squares. For each vertex $x \in V(\mathbf{P}_q)$, its neighbors are $x + z^0, x + z^2, x + z^4, \dots, x + z^{q-3}$, and we now take this (cyclic) sequence to define the local rotation around x . This defines an orientable embedding Π_q of \mathbf{P}_q which is known as the *Paley map*. It has the following properties (see [12]):

Theorem 3.2 *Suppose that $q \equiv 1 \pmod{8}$. Then the Paley map Π_q is self-dual – the geometric dual graph \mathbf{P}_q^* is isomorphic to \mathbf{P}_q . Its genus is equal to $(q^2 - 9q + 8)/8$. It has q faces, each of length $(q - 1)/2$ and bounded by a cycle of \mathbf{P}_q .*

We refer to [12] for more details concerning this interesting map.

From now on we assume that $q \equiv 1 \pmod{8}$. Let T_q be the triangulation of the orientable surface of genus $(q^2 - 9q + 8)/8$ obtained from the Paley map Π_q by adding a new vertex into each face and joining it to all vertices on the boundary of that face. Let $V_q = V(\mathbf{P}_q)$ and let V_q^* be the added vertices. Then $|V_q| = |V_q^*| = q$.

Theorem 3.3 *Let $q \equiv 1 \pmod{8}$ be a prime power. The triangulation T_q contains no subdivision of the complete graph of order $\geq \lambda\sqrt{q}$, where $\lambda = \frac{1}{4}(2 + \sqrt{198}) < 4.0178$.*

Proof. Let $s = \sqrt{q}$ and $k = \lceil \lambda s \rceil$. Suppose that T_q contains a subdivision K of K_k . Let $B_0 \subseteq V(K)$ be the set of branch vertices, and let $b = |B_0 \cap V_q|$ and $b^* = |B_0 \cap V_q^*|$. Clearly, $b + b^* = k$.

Let us count the number of subdivided edges in K . Since no two vertices in V_q^* are adjacent, $B_0 \cap V_q^*$ gives rise to $\binom{b^*}{2}$ subdivided edges. Concerning the set $B_0 \cap V_q$, Theorem 3.1 shows that this set gives rise to at least $\frac{1}{2} \binom{b}{2} - \frac{b(q-b)}{4s}$ subdivided edges. Since the total number of vertices of T_q is equal to $2q$, we see that:

$$k + \binom{b^*}{2} + \frac{1}{2} \binom{b}{2} - \frac{b(q-b)}{4s} \leq 2q. \quad (1)$$

Let $b^* = \alpha^* s$ and $b = \alpha s$. Then (1) expands to the following condition

$$s(2\alpha^{*2} + \alpha^2 - \alpha - 8) + 2\alpha^* + 3\alpha + \alpha^2 \leq 0. \quad (2)$$

Let $\Lambda = \alpha^* + \alpha$. Then (2) implies that

$$2(\Lambda - \alpha)^2 + \alpha^2 - \alpha - 8 < 0. \quad (3)$$

This quadratic inequality has no solutions if

$$8\Lambda^2 - 8\Lambda - 97 \geq 0. \quad (4)$$

Since $\Lambda s = k = \lceil \lambda s \rceil \geq \lambda s$, we have $\Lambda \geq \lambda = \frac{1}{4}(2 + \sqrt{198})$. This implies (4) and proves that (3) has no solution. This contradiction completes the proof. \square

Theorem 3.3 now implies:

Theorem 3.4 *Let $q \equiv 1 \pmod{8}$ be a prime power. If the chromatic number of \mathbf{P}_q is at least $\lambda\sqrt{q}$, where $\lambda = \frac{1}{4}(2 + \sqrt{198})$, then the triangulation T_q fails to satisfy the Hajós Conjecture.*

In the above theorem, it suffices to assume that $\chi(T_q) \geq \lambda\sqrt{q}$. But since the vertex set V_q^* is independent in T_q , $\chi(T_q) \leq \chi(\mathbf{P}_q) + 1$. Therefore, we find it more pleasing to use the natural value $\chi(\mathbf{P}_q)$.

The independence numbers $\alpha(\mathbf{P}_q)$ of Paley graphs of prime order q less than 7000 were computed by Shearer [8]. Using these calculations and estimate the chromatic number by $\chi(\mathbf{P}_q) \geq \lceil q/\alpha(\mathbf{P}_q) \rceil$, one finds out that Theorem 3.4 can be applied for many values of q (and when q gets large, almost all of them are good). All such primes $q \leq 7000$ with $q \equiv 1 \pmod{8}$ are collected in Table 1.

As stated after Theorem 3.1, it is believed that for every prime q , $\omega(\mathbf{P}_q) = o(\sqrt{q})$. If true, this yields infinitely many cases where Theorem 3.4 can be applied.

q	$\alpha(\mathbf{P}_q)$	$\lceil q/\alpha(\mathbf{P}_q) \rceil$	$\lceil \lambda\sqrt{q} \rceil$	q	$\alpha(\mathbf{P}_q)$	$\lceil q/\alpha(\mathbf{P}_q) \rceil$	$\lceil \lambda\sqrt{q} \rceil$
3697	15	247	245	5953	17	351	310
4217	15	282	261	6073	19	320	314
4441	15	297	268	6089	17	359	314
4457	15	298	269	6113	19	322	315
4649	17	274	274	6121	18	341	315
4673	17	275	275	6217	19	328	317
4721	17	278	277	6257	19	330	318
4729	17	279	277	6329	19	334	320
4793	16	300	279	6337	17	373	320
4817	17	284	279	6353	17	374	321
4937	17	291	283	6361	18	354	321
4969	17	293	284	6449	19	340	323
4993	17	294	284	6473	19	341	324
5081	17	299	287	6481	17	382	324
5153	17	304	289	6529	17	385	325
5233	17	308	291	6553	19	345	326
5281	17	311	292	6569	19	346	326
5297	17	312	293	6577	19	347	326
5393	17	318	296	6673	19	352	329
5441	17	321	297	6689	19	353	329
5449	17	321	297	6737	18	375	330
5569	18	310	300	6761	20	339	331
5657	18	315	303	6793	19	358	332
5737	17	338	305	6833	19	360	333
5801	17	342	307	6841	17	403	333
5849	17	345	308	6857	17	404	333
5857	17	345	308	6961	19	367	336
5881	19	310	309	6977	18	388	336
5897	17	347	309				

Table 1: Some good values of q

4 Conclusions

We have proved that there is a variety of reasons why a triangulation of some surface may fail to satisfy the Hajós Conjecture. In this sense, this note can be viewed as an echo to the stimulating work of Thomassen [11]. As noted in that paper, graphs embedded in a surface Σ with sufficiently large edge-

width satisfy the Hajós Conjecture. Recall that the *edge-width* of a graph embedded in a nonplanar surface is the length of a shortest noncontractible cycle. In the above observation of Thomassen, the required width depends on Σ . However, the following strengthening may be true:

Conjecture 4.1 *There is an absolute constant w_0 such that every graph which can be embedded in some surface with edge-width at least w_0 satisfies the Hajós Conjecture.*

Kühn and Osthus [5] proved that graphs whose girth is at least 186 satisfy the Hajós conjecture. This excludes the most obvious possibility of counterexamples to Conjecture 4.1. (Note that the edge-width of a graph of girth g is at least g .)

We do not dare to estimate what the best possible value for w_0 may be. However, if we restrict ourselves to triangulations, no counterexamples of edge-width 4 are known.

Problem 4.2 *Does every triangulation without noncontractible triangles satisfy the Hajós Conjecture?*

For a graph H and a positive integer t , let \tilde{H}_t be the graph obtained from the union of H and the clique K_t by adding all edges between them. Clearly, the Hajós Conjecture holds for H if and only if it holds for \tilde{H}_t .

Let v and e (\tilde{v} and \tilde{e}) be the number of vertices and edges of H (and \tilde{H}_t), respectively. Clearly, $\tilde{v} = v + t$ and $\tilde{e} = e + \binom{t}{2} + vt$. If \tilde{H}_t triangulates some surface, then \tilde{e} is divisible by 3. This holds if and only if one of the following holds:

- (A) $e \equiv 0 \pmod{3}$ and $t \equiv 0 \pmod{3}$,
- (B) $v + e \equiv 0 \pmod{3}$ and $t \equiv 1 \pmod{3}$, or
- (C) $e - v \equiv 2 \pmod{3}$ and $t \equiv 2 \pmod{3}$.

Conjecture 4.3 *For every graph H , there exists a constant t_0 such that for every $t \geq t_0$ for which one of (A)–(C) is satisfied, the graph \tilde{H}_t triangulates some surface.*

Conjecture 4.3 would imply that for every counterexample H to the Hajós Conjecture for which either $e \equiv 0 \pmod{3}$, $v + e \equiv 0 \pmod{3}$, or $e - v \equiv 2 \pmod{3}$, there are infinitely many triangulations of the form \tilde{H}_t which violate the Hajós Conjecture.

Conjecture 4.3 would follow from a solution of a more general open problem which has been raised in [7, Problem 4.4.10]:

Problem 4.4 *Is there an $\varepsilon > 0$ such that every graph of order n , with e edges, where e is divisible by 3, and with minimum degree at least $(1 - \varepsilon)n$ triangulates some surface?*

A positive answer to Problem 4.4 would imply that almost one third of random graphs in $\mathcal{G}(n, p)$, where $p > 1 - \varepsilon$, would be graphs of some triangulations. As shown by Erdős and Fajtlowicz [4], almost all of these graphs would also be counterexamples to Conjecture 1.1.

References

- [1] B. Bollobás, Random graphs, Second ed., Cambridge University Press, Cambridge, 2001.
- [2] P. Catlin, Hajós' graph-coloring conjecture: variations and counterexamples, J. Combin. Theory, Ser. B 26 (1979) 268–274.
- [3] G. A. Dirac, A property of 4-chromatic graphs and some remarks on critical graphs, J. London Math. Soc. 27 (1952) 85–92.
- [4] P. Erdős, S. Fajtlowicz, On the conjecture of Hajós, Combinatorica 1 (1981) 141–143.
- [5] D. Kühn, D. Osthus, Topological minors in graphs of large girth, J. Combin. Theory, Ser. B 86 (2002) 364–380.
- [6] W. Mader, $3n - 5$ edges do force a subdivision of K_5 , Combinatorica 18 (1998) 569–595.
- [7] B. Mohar and C. Thomassen, Graphs on Surfaces, Johns Hopkins University Press, Baltimore, 2001.
- [8] J. B. Shearer,
<http://www.research.ibm.com/people/s/shearer/indpal.html>
- [9] A. G. Thomason, Paley graphs and Weil's theorem, Talk at the British Combinatorial Conference, Southampton, 1983 (unpublished).
- [10] C. Thomassen, Five-coloring graphs on the torus, J. Combin. Theory Ser. B 62 (1994) 11–33.
- [11] C. Thomassen, Some remarks on Hajós' conjecture, J. Combin. Theory, Ser. B 93 (2005) 95–105.
- [12] A. T. White, Graphs of groups on surfaces. Interactions and models, North-Holland, Amsterdam, 2001.