

UNIVERSITY OF LJUBLJANA  
INSTITUTE OF MATHEMATICS, PHYSICS AND MECHANICS  
DEPARTMENT OF MATHEMATICS  
JADRANSKA 19, 1 111 LJUBLJANA, SLOVENIA

**Preprint series, Vol. 44 (2006), 1005**

CIRCULAR COLOURING THE  
PLANE

Matt Devos      Javad Ebrahimi  
Mohammad Ghebleh  
Luis Goddyn , Bojan Mohar  
Reza Naserasr

ISSN 1318-4865

July 5, 2006

Ljubljana, July 5, 2006

# CIRCULAR COLOURING THE PLANE

MATT DEVOS , JAVAD EBRAHIMI , MOHAMMAD GHEBLEH , LUIS GODDYN ,  
BOJAN MOHAR <sup>\*</sup>, AND REZA NASERASR <sup>†</sup>

**Abstract.** The *unit distance graph*  $\mathcal{R}$  is the graph with vertex set  $\mathbb{R}^2$  in which two vertices (points in the plane) are adjacent if and only if they are at Euclidean distance 1. We prove that the circular chromatic number of  $\mathcal{R}$  is at least 4, thus “tightening” the known lower bound on the chromatic number of  $\mathcal{R}$ .

**Key words.** graph colouring, circular colouring, unit distance graph

**AMS subject classifications.** 05C15, 05C10, 05C62

**1. Introduction.** The *unit distance graph*  $\mathcal{R}$  is defined to be the graph with vertex set  $\mathbb{R}^2$  in which two vertices (points in the plane) are adjacent if and only if they are at Euclidean distance 1. Every subgraph of  $\mathcal{R}$  is also said to be a *unit distance graph*. It is known that (cf. [1, 2])

$$4 \leq \chi(\mathcal{R}) \leq 7,$$

and that (cf. [3, pp. 59–65])

$$\frac{32}{9} \leq \chi_f(\mathcal{R}) \leq 4.36.$$

Here  $\chi(\mathcal{R})$  and  $\chi_f(\mathcal{R})$  denote the chromatic number and the fractional chromatic number of  $\mathcal{R}$ , respectively. In this paper we study the circular chromatic number of the unit distance graph  $\mathcal{R}$ .

Let  $r \geq 2$ ,  $a, b \in [0, r)$ , and  $a \leq b$ . We define the *circular distance* of  $a$  and  $b$ , denoted by  $\delta(a, b) = \delta_r(a, b)$ , to be  $\min\{b - a, r + a - b\}$ . One may identify the interval  $[0, r)$  with a circle  $C^r$  with perimeter  $r$  and then  $\delta(a, b)$  will be the distance between  $a$  and  $b$  in  $C^r$ .

If  $a, b \in [0, r)$  (or equivalently  $a, b \in C^r$ ), we define the *circular interval from  $a$  to  $b$* , denoted  $[a, b]$ , as follows (see Figure 1.1):

$$[a, b] = \begin{cases} \{x \mid a \leq x \leq b\} & \text{if } a \leq b, \\ \{x \mid 0 \leq x \leq b \text{ or } a \leq x < r\} & \text{if } a > b. \end{cases}$$

An  *$r$ -circular colouring* of a graph  $G$ , is a function  $c : V(G) \rightarrow C^r$  such that for every edge  $xy$  in  $G$ ,  $\delta(c(x), c(y)) \geq 1$ . The *circular chromatic number* of  $G$ , denoted by  $\chi_c(G)$ , is

$$\chi_c(G) = \inf\{r \mid G \text{ admits an } r\text{-circular colouring}\}.$$



It is well known [4] that for every graph  $G$ ,  $\chi_f(G) \leq \chi_c(G) \leq \chi(G)$ . For the unit distance graph  $\mathcal{R}$ , these inequalities give

$$\frac{32}{9} \leq \chi_f(\mathcal{R}) \leq \chi_c(\mathcal{R}) \leq \chi(\mathcal{R}) \leq 7.$$

---

<sup>\*</sup>Department of Mathematics, Simon Fraser University, Burnaby, British Columbia, V5A 1S6, Canada.

<sup>†</sup>Department of Combinatorics and Optimization, Faculty of Mathematics, University of Waterloo, Waterloo, Ontario, N2L 3G1, Canada.

FIG. 1.1. *Circular intervals (clockwise direction is the positive direction)*FIG. 2.1. *The unit distance graph  $H_{a,b}$* 

We improve the lower bound for  $\chi_c(\mathcal{R})$  to 4. We give two proofs of this result. The second one is constructive and gives a construction of finite unit distance graphs with circular chromatic number arbitrarily close to 4.

**2. Proof.** Let  $a$  and  $b$  be two points in the plane and let  $d(a,b)$  denote the Euclidean distance between  $a$  and  $b$ . If  $d(a,b) = \sqrt{3}$ , then we may find points  $x$  and  $y$  in the plane such that the subgraph of  $\mathcal{R}$  induced on the set  $\{a, b, x, y\}$  is isomorphic to the graph  $H$  obtained by deleting one edge from  $K_4$  (see Figure 2.1). We denote this unit distance graph by  $H_{a,b}$ . On the other hand, it is easy to see that in any embedding of  $H$  as a unit distance graph in the plane, the Euclidean distance between the two vertices of degree 2 in  $H$  is  $\sqrt{3}$ .

**LEMMA 2.1.** *Let  $0 < \varepsilon < 1$  and  $a, b \in \mathbb{R}^2$  with  $d(a,b) = \sqrt{3}$ . Let  $c$  be a  $(3 + \varepsilon)$ -circular colouring of  $H_{a,b}$ . Then  $\delta(c(a), c(b)) \leq \varepsilon$ .*

*Proof.* Without loss of generality, we may assume  $c(a) = 0$ . Since  $a, x, y$  form a triangle in  $H_{a,b}$ , we have  $c(x) \in [1, 1 + \varepsilon]$  and  $c(y) \in [2, 2 + \varepsilon]$  up to symmetry. On the other hand,  $b$  is adjacent to both  $x$  and  $y$ . Thus

$$\begin{aligned} c(b) &\in [c(x) + 1, c(x) - 1] \cap [c(y) + 1, c(y) - 1] \\ &\subseteq [2, \varepsilon] \cap [-\varepsilon, 1 + \varepsilon] \\ &= [-\varepsilon, \varepsilon]. \end{aligned}$$

The last equality is true since  $1 + \varepsilon < 2$ .  $\square$

**THEOREM 2.2.**  $\chi_c(\mathcal{R}) \geq 4$ .

*Proof.* Suppose that  $c$  is a  $(3 + \varepsilon)$ -circular colouring of  $\mathcal{R}$  where  $0 \leq \varepsilon < 1$ . Let

$$\mu = \sup\{\delta(c(a), c(b)) \mid a, b \in \mathbb{R}^2 \text{ and } d(a,b) = \sqrt{3}\}.$$

By Lemma 2.1,  $\mu \leq \varepsilon$ . By the definition of  $\mu$ , for every  $0 < \mu' < \mu$ , there exist points  $a$  and  $b$  at distance  $\sqrt{3}$  in the plane such that  $\delta(c(a), c(b)) > \mu'$ . Consider the graph

$H_{a,b}$  as in Figure 2.1. Without loss of generality we may assume

$$0 = c(a) \leq c(b) < c(x) < c(y) \leq 2 + \varepsilon.$$

Since  $3 + \varepsilon < 4$ , we have

$$\delta(c(a), c(x)) = c(x) = \delta(c(a), c(b)) + \delta(c(b), c(x)) > \mu' + 1.$$

On the other hand since  $a$  and  $x$  are at distance 1, there exists a point  $z$  which is at distance  $\sqrt{3}$  from both  $a$  and  $x$ . Therefore

$$1 \leq \delta(c(a), c(x)) \leq \delta(c(a), c(z)) + \delta(c(z), c(x)) \leq 2\mu.$$

Hence  $1 + \mu' < 2\mu$  and since this is true for every  $\mu' < \mu$ , we have  $\mu \geq 1$ . This is a contradiction since  $\mu \leq \varepsilon < 1$ .  $\square$

**3. A constructive proof.** The graph  $G_0 = K_2$  is obviously a unit distance graph. In our construction of graphs  $G_n$  ( $n \geq 0$ ) we distinguish two vertices in each of them. To emphasize the distinguished vertices  $x$  and  $y$  of  $G_n$ , we write  $G_n^{x,y}$ . We identify subgraphs of  $\mathcal{R}$  with their geometric representation given by their vertex set.

For  $n \geq 0$ , the graph  $G_{n+1}$  is constructed recursively from four copies of  $G_n$ . Let  $S = V(G_n^{x,y}) \subseteq \mathbb{R}^2$ . Let us rotate the set  $S$  in the plane about the point  $x$ , so that the image  $y'$  of  $y$  under this rotation is at distance 1 from  $y$ . Let  $S'$  be the image of  $S$  under this rotation. Let  $T$  be the set of all points in  $S \cup S'$  and their reflections across the line  $yy'$ . In particular let  $z \in T$  be the reflection of  $x$  across the line  $yy'$ . We define  $G_{n+1}^{x,z}$  to be the subgraph of  $\mathcal{R}$  induced on  $T$ . This construction is depicted in Figure 3.1.




FIG. 3.1. Construction of  $G_{n+1}$  from  $G_n$

**LEMMA 3.1.** *For every  $n \geq 1$ ,  $\chi_c(G_n) \geq 4 - 2^{1-n}$ . Moreover, for every  $r = 4 - 2^{1-n} + \varepsilon$  with  $0 \leq \varepsilon < 2^{1-n}$ , and every circular  $r$ -colouring  $c$  of  $G_n^{x,z}$ , we have  $\delta(c(x), c(z)) \leq 2^{n-1}\varepsilon$ .*

*Proof.* We use induction on  $n$ . The case  $n = 1$  is proved in Lemma 2.1. Let  $n \geq 1$  and  $G_{n+1}^{x,z}$  be as shown in Figure 3.1. Let  $r = 4 - 2^{1-n} + \varepsilon$  for some  $\varepsilon \geq 0$  and let  $c$  be a circular  $r$ -colouring of  $G_{n+1}$ . Without loss of generality we may assume that  $c(x) = 0$ . By the induction hypothesis,  $\delta(0, c(y))$  and  $\delta(0, c(y'))$  are both at most  $2^{n-1}\varepsilon$ . Hence  $\delta(c(y), c(y')) \leq 2^n\varepsilon$ . On the other hand, since  $y$  and  $y'$  are adjacent in  $G_{n+1}$ , we have  $\delta(c(y), c(y')) \geq 1$ . Therefore  $\varepsilon \geq 2^{-n}$  and we have  $\chi_c(G_{n+1}) \geq 4 - 2^{1-n} + 2^{-n} = 4 - 2^{-n}$ .

Now let  $r = 4 - 2^{-n} + \varepsilon$  for some  $0 \leq \varepsilon < 2^{-n}$ , and let  $c$  be a circular  $r$ -colouring of  $G_{n+1}$  with  $c(x) = 0$ . Note that  $r = 4 - 2^{1-n} + \varepsilon'$  with  $\varepsilon' = 2^{-n} + \varepsilon < 2^{1-n}$ . By

the induction hypothesis,  $\delta(0, c(y))$ ,  $\delta(0, c(y'))$ ,  $\delta(c(z), c(y))$  and  $\delta(c(z), c(y'))$  are all at most  $2^{n-1}\varepsilon' < 1$ . Therefore we have

$$c(y), c(y') \in [-2^{n-1}\varepsilon', 2^{n-1}\varepsilon']$$

and

$$c(z) \in [c(y) - 2^{n-1}\varepsilon', c(y) + 2^{n-1}\varepsilon'] \cap [c(y') - 2^{n-1}\varepsilon', c(y') + 2^{n-1}\varepsilon'].$$

Since  $\delta(c(y), c(y')) \geq 1$ , one of  $c(y)$  and  $c(y')$ , say  $c(y)$ , is in the circular interval  $[-2^{n-1}\varepsilon', 2^{n-1}\varepsilon' - 1]$ , and  $c(y') \in [-2^{n-1}\varepsilon' + 1, 2^{n-1}\varepsilon']$ . Therefore

$$[c(y) - 2^{n-1}\varepsilon', c(y) + 2^{n-1}\varepsilon'] \subseteq [-2^n\varepsilon', 2^n\varepsilon' - 1] = [-2^n\varepsilon', 2^n\varepsilon]$$

and

$$[c(y') - 2^{n-1}\varepsilon', c(y') + 2^{n-1}\varepsilon'] \subseteq [-2^n\varepsilon' + 1, 2^n\varepsilon'] = [-2^n\varepsilon, 2^n\varepsilon'].$$

Finally, since  $\varepsilon' < 2^{1-n}$ , we have  $2^n\varepsilon' < r - 2^n\varepsilon'$ . Hence

$$c(z) \in [-2^n\varepsilon', 2^n\varepsilon] \cap [-2^n\varepsilon, 2^n\varepsilon'] = [-2^n\varepsilon, 2^n\varepsilon].$$

This completes the induction step.  $\square$

Let us observe that, when constructing  $G_{n+1}$  from four copies of  $G_n$ , it may happen that vertices in distinct copies of  $G_n$  correspond to the same points in the plane. Additionally, it may happen that some edges between vertices in distinct copies of  $G_n$  are introduced. We may define in the same way a sequence of abstract graphs  $H_n$ , where none of these two issues occur. Clearly  $\chi_c(G_n) \geq \chi_c(H_n)$ , but we cannot argue equality in general. The proof of Lemma 3.1 applied to the graphs  $H_n$  gives slightly more:

**THEOREM 3.2.** *For every  $n \geq 0$ ,  $\chi_c(H_n) = 4 - 2^{1-n}$ .*

*Proof.* The cases  $n = 0, 1$  are trivial. Let  $n \geq 1$  and let  $H_{n+1}$  be as in Figure 3.1. Let  $r = 4 - 2^{-n} = 4 - 2^{1-n} + 2^n$ . By the proof of Lemma 3.1,  $H_n^{x,y}$  admits a circular  $r$ -colouring  $c_1$  with  $c_1(x) = 0$  and  $c_1(y) = \frac{1}{2}$ . Similarly the graphs  $H_n^{x,y'}$ ,  $H_n^{y,z}$  and  $H_n^{y',z}$  admit circular  $r$ -colourings  $c_2$ ,  $c_3$  and  $c_4$ , respectively, with  $c_2(x) = 0$ ,  $c_2(y') = c_4(y') = -\frac{1}{2}$ ,  $c_3(y) = \frac{1}{2}$ , and  $c_3(z) = c_4(z) = 0$ . Now a circular  $r$ -colouring  $c$  of  $H_{n+1}$  can be obtained by combining the partial colourings  $c_1, c_2, c_3, c_4$ .  $\square$

The construction of this section gives an infinite subgraph of  $\mathcal{R}$  with circular chromatic number at least 4. It remains open whether or not  $\mathcal{R}$  has a finite subgraph with the same property.

#### REFERENCES

- [1] H. HADWIGER AND H. DEBRUNNER, *Combinatorial geometry in the plane*, Holt, Rinehart and Winston, New York, 1964.
- [2] L. MOSER AND W. MOSER, *Solution to problem 10*, Canad. Math. Bull. 4 (1961), 187–189.
- [3] E. R. SCHEINERMAN AND D. H. ULLMAN, *Fractional graph theory*, John Wiley & Sons Inc., New York, 1997.
- [4] X. ZHU, *Circular chromatic number: a survey*, Discrete Math. 229 (2001), 371–410.