

UNIVERSITY OF LJUBLJANA  
INSTITUTE OF MATHEMATICS, PHYSICS AND MECHANICS  
DEPARTMENT OF MATHEMATICS  
JADRANSKA 19, 1000 LJUBLJANA, SLOVENIA

**Preprint series, Vol. 46 (2008), 1044**

BACKBONE COLORINGS AND  
GENERALIZED MYCIELSKI'S  
GRAPH

Jozef Miškuf      Riste Škrekovski  
Martin Tancer

ISSN 1318-4865

March 10, 2008

Ljubljana, March 10, 2008

# Backbone Colorings and Generalized Mycielski's Graphs\*

Jozef Miškuf<sup>†</sup>     Riste Škrekovski<sup>‡</sup>     Martin Tancer<sup>§</sup>

## Abstract

For a graph  $G$  and its spanning tree  $T$  the *backbone chromatic number*,  $\text{BBC}(G, T)$ , is defined as the minimum  $k$  such that there exists a coloring  $c: V(G) \rightarrow \{1, 2, \dots, k\}$  satisfying  $|c(u) - c(v)| \geq 1$  if  $uv \in E(G)$  and  $|c(u) - c(v)| \geq 2$  if  $uv \in E(T)$ .

Broersma et al. [1] asked whether there exists a constant  $c$  such that for every triangle-free graph  $G$  with an arbitrary spanning tree  $T$  the inequality  $\text{BBC}(G, T) \leq \chi(G) + c$  holds. We answer this question negatively by showing the existence of triangle-free graphs  $R_n$  and their spanning trees  $T_n$  such that  $\text{BBC}(R_n, T_n) = 2\chi(R_n) - 1 = 2n - 1$ .

In order to answer the question we obtain a result of independent interest. We modify the well known Mycielski's construction and construct triangle-free graphs  $J_n$ , for every integer  $n$ , with chromatic number  $n$  and 2-tuple chromatic number  $2n$  (here 2 can be replaced by any integer  $t$ ).

**Keywords:** backbone coloring, graph coloring, generalized Mycielski's construction, triangle-free graph.

**MSC:** 05C15

---

\*This work was partially supported by the Czech-Slovenian bilateral research project 15/2006-2007 and by the Slovak-Slovenian bilateral research project 1/2006-2007.

<sup>†</sup>Institute of Mathematics, Faculty of Science, University of Pavol Jozef Šafárik, Jesenná 5, 041 54 Košice, Slovakia. E-mail: [jozef.miskuf@upjs.sk](mailto:jozef.miskuf@upjs.sk)

<sup>‡</sup>Department of Mathematics, Faculty of Mathematics and Physics, University of Ljubljana, Jadranska 19, 1000 Ljubljana, Slovenia. Supported in part by ARRS Research Program P1-0297.

<sup>§</sup>Institute for Theoretical Computer Science (supported by project 1M0545 of The Ministry of Education of the Czech Republic), Faculty of Mathematics and Physics, Charles University, Malostranské nám. 25, 118 00 Prague, Czech Republic. E-mail: [tancer@kam.mff.cuni.cz](mailto:tancer@kam.mff.cuni.cz)

# 1 Introduction

## 1.1 Backbone colorings

The backbone coloring problem is related to frequency assignment problems in the following way, the transmitters are represented by the vertices of a graph and they are adjacent in the graph if the corresponding transmitters are close enough or transmitters are strong enough. The problem is to assign frequency channels to the transmitters in such a way that the interference is kept at an "acceptable" level. One way of putting these requirements together is following: Given graphs  $G_1, G_2$  such that  $G_1$  is a spanning subgraph of  $G_2$ . Determine a coloring of  $G_2$  that satisfies certain restriction of one type in  $G_1$  and of the other type in  $G_2$ .

In this concept, backbone colorings were introduced and motivated and put into a general framework of related coloring problems in [1]. Let us recall some basic definitions. In the sequel we deal with undirected simple graphs, i.e. without loops and/or multiedges. By the symbol  $[n]$  we understand the set  $\{1, 2, \dots, n\}$ , by the symbol  $\chi(G)$  the chromatic number of  $G$ , and by the symbol  $G[W]$  the subgraph induced by the vertex set  $W \subseteq V(G)$ . For a graph  $G$ , we define a coloring  $\nu : V \rightarrow \{1, 2, \dots, k\}$  to be a *backbone  $k$ -coloring* of a graph  $G$  with a backbone graph  $H \subseteq G$ , if for every two different vertices  $u$  and  $v$  of  $G$ , it holds

- $|\nu(u) - \nu(v)| \geq 1$ , if  $uv \in E(G) \setminus E(H)$ ;
- $|\nu(u) - \nu(v)| \geq 2$ , if  $uv \in E(H)$ .

The minimum  $k$  for which  $G$  with backbone  $H$  admits a backbone  $k$ -coloring is called the *backbone chromatic number* of  $G$  with backbone  $H$ . It is denoted by  $\text{BBC}(G, H)$ . In this paper we consider only the case when a backbone graph  $H$  is acyclic.

We refer to several results concerning backbone colorings of graphs. The connection between the backbone chromatic number and the chromatic number is studied in [1]. The authors showed that the backbone chromatic number of a graph  $G$  is at most  $2\chi(G) - 1$ , while they provided examples where this bound is attained. To show this inequality it is sufficient to color the graph  $G$  with colors  $1, 3, \dots, 2\chi(G) - 1$ . The decision problem if there exists a backbone coloring of a graph  $G$  with backbone tree  $T$  with  $l$  colors is NP-complete for  $l \geq 5$ . Broersma et al. in [3] showed that the backbone chromatic number of planar graphs with backbone matchings is at most six. Other results on backbone colorings appear in [2, 4, 9].

We deal with the intriguing question posed by Broersma et al. [1].

**Question 1.1.** *Does there exist a constant  $c$  such that  $\text{BBC}(G, T) \leq \chi(G) + c$  holds for every triangle-free graphs  $G$  with  $T$  being a tree?*

We will present infinite class of triangle-free graphs answering the question negatively. More precisely, for every integer  $n$ , we will show the existence of a triangle-free graph  $G$  with a backbone tree  $T$  such that  $\text{BBC}(G, T) = 2\chi(G) - 1 = 2n - 1$ .

## 1.2 Triangle-free graphs and their colorings

For integers  $k$  and  $t$  we define a  $t$ -tuple  $k$ -coloring of a graph  $G$  to be a function  $c: V(G) \rightarrow \binom{[k]}{t}$  such that  $c(u) \cap c(v) = \emptyset$  whenever  $uv \in E(G)$ . The minimum possible  $k$  that  $G$  has a  $t$ -tuple  $k$ -coloring is called the  $t$ -tuple chromatic number, denoted by  $\chi_t(G)$ .

The procedure of giving a negative answer to Question 1.1 will proceed in the following steps:

- Step I. For a given triangle-free graph  $G$ , we will construct an infinite triangle-free graph  $R_G$  with a backbone tree  $T_G$  such that  $\text{BBC}(R_G, T_G) \geq \chi_2(G) - 1$  and  $\chi(R_G) = \chi(G)$ .
- Step II. For a given triangle-free graph  $G$ , we will present a Mycielski-type construction of a triangle-free graph  $J(G)$  such that  $\chi_2(J(G)) \geq \chi_2(G) + 2$  and  $\chi(J(G)) \leq \chi(G) + 1$ . In particular, it follows that  $\chi(J_n) = n$  and  $\chi_2(J_n) = 2n$ , where  $J_n = J^{n-2}(K_2)$  and  $K_2$  is the complete graph on 2 vertices.
- Step III. From the previous two steps,  $\text{BBC}(R_{J_n}, T_{J_n}) \geq 2n - 1 = 2\chi(R_{J_n}) - 1$ . The graph  $R_{J_n}$  is infinite; however, by the principle of compactness there exists a finite (connected) subgraph  $R_n \subseteq R_{J_n}$  such that  $\text{BBC}(R_n, T_{J_n}[V(R_n)]) \geq 2n - 1$ .  $T_{J_n}[V(R_n)]$  is a subforest of  $R_n$  thus it can be extended to a spanning tree  $T_n$  of  $R_n$ . We know that  $\text{BBC}(R_n, T_n) \geq 2n - 1$  and  $\chi(R_n) \leq \chi(R_{J_n}) = n$ . Actually, equalities hold since  $\text{BBC}(G, T) \leq 2\chi(G) - 1$  for any graph  $G$  with backbone  $T$ .

The construction from Step I follows an idea of Broersma et al. [1]. It will be described in Section 2.

The crucial step is Step II. Its task is to construct a triangle-free graph  $J_n$ , for every integer  $n$ , such that  $\chi(J_n) = n$  and  $\chi_2(J_n) = 2n$ . This construction may be of an independent interest. The well known *fractional chromatic number* of a graph  $G$  is defined as

$$\chi_f = \inf_{t \in \mathbb{N}} \frac{\chi_t(G)}{t}.$$

Since  $\chi(G) \geq \frac{\chi_2(G)}{2} \geq \chi_f(G)$ , it would be natural to look for a class, say  $F_n$ , of triangle-free graphs such that  $\chi(F_n) = \chi_f(F_n) = n$ . Unfortunately, we are not aware of such a class of graphs. There is a lot of known examples of triangle-free

graphs with a large chromatic number, however it seems that none of them is the case. We just give a note on some of them:

Larsen, Propp and Ullman [7] proved that a Mycielski graph with the chromatic number  $n$  has fractional chromatic number approximately equal to  $\sqrt{2n}$ .

Kneser graphs  $KG_{3n-1,n}$  are triangle-free graphs with chromatic number  $n+1$  and fractional chromatic number  $\frac{3n-1}{n} < 3$ . More details about colorings of Kneser graphs can be found e.g. in [6, 8].

By the probabilistic method, Erdős [5] showed that exist triangle-free graphs with an arbitrary large chromatic number actually proving that the independence number of such graphs is small (and thus even fractional chromatic number is large). However, it does not give an intuition, how far away the chromatic number and the fractional chromatic number are.

The construction of graphs  $J_n$  is a generalization of Mycielski's construction. It will be precisely described in Section 3.

For Step III we just remark that with some extra effort it is possible to work just with finite graphs and avoid any use of the principle of compactness. However, some of the proofs would be more complicated and more technical, thus we rather prefer to work with infinite graphs.

## 2 Relation between backbone colorings and 2-tuple colorings

In this section, for a given graph  $G$  we construct a pair of graphs  $(R_G, T_G)$  as mentioned in Step I in the introduction.

**Definition 2.1.** Let  $G$  be a graph. We define a pair of infinite graphs  $(R_G, T_G)$  in the following way:

1. The graph  $R_G$  is the OR-product of  $G$  and  $\mathbb{N}$  (as independent set), i.e.
  - $V(R_G) = V(G) \times \mathbb{N}$ ;
  - $E(R_G) = \{(v_1, n_1), (v_2, n_2)\} \mid v_1 v_2 \in E(G)\}$ .
2.  $T_G$  is a spanning tree of  $G$  defined recursively in the following way: Fix a good linear ordering  $\preceq$  on  $V(R_G)$ . Let  $T_G^1$  is the graph with just one vertex – the smallest element of  $V(R_G)$ . Suppose that  $T_G^i$  is already defined. Let us define  $T_G^{i+1} \supset T_G^i$ : First, we call *occupied* all the vertices of  $T_G^i$ . Gradually for every vertex  $(v, k) \in T_G$  (from the smallest one to the largest one) and for every edge  $uv \in E(G)$ , choose  $j$  such that  $(u, j)$  is the smallest possible unoccupied vertex and add this vertex among the occupied vertices. Moreover, add the vertex  $(u, j)$  among the vertices of  $T_G^{i+1}$  and the edge  $\{(v, k), (u, j)\}$  among the edges of  $T_G^{i+1}$ . Finally, we define  $T_G = \bigcup_{i \in \mathbb{N}} T_G^i$ .




Figure 1: The path  $P_3$  and the pair  $(R_{P_3}, T_{P_3})$ , where the ordering  $\preceq$  is chosen so that vertices with higher altitude in the figure are smaller in  $\preceq$ . Labels at vertices are the smallest such  $i$  that the corresponding vertex belongs to  $T_{P_3}^i$ .

An example of the construction is depicted on Figure 1. Notice, that the spanning tree  $T_G$  is not defined uniquely; however, for our purposes it is not important to have a unique definition. The following lemma easily follows from the construction.

**Lemma 2.2.** *For any graph  $G$ , the pair  $(R_G, T_G)$  has the following properties:*

1.  $T_G$  is a spanning tree of  $R_G$ .
2. If  $G$  is triangle-free then  $R_G$  is triangle-free.
3. For every vertex  $(v, j)$  of  $R_G$  and for every edge  $uv$  of  $G$ , there exists an integer  $j'$  such that  $\{(v, j), (u, j')\}$  is an edge of  $T_G$ .

□

The following proposition relates the 2-tuple chromatic number of a graph  $G$  and the backbone chromatic number of the pair  $(R_G, T_G)$ .

**Proposition 2.3.** *Let  $G$  be a graph. Then*

1.  $\chi(R_G) = \chi(G)$ ;
2.  $\text{BBC}(R_G, T_G) \geq \chi_2(G) - 1$ .

*Proof.* We prove each of the claims separately:

1. The graph  $R_G[V(G) \times \{1\}]$  is isomorphic to  $G$ , hence  $\chi(G) \leq \chi(R_G)$ .

On the other hand, any coloring of  $G$  induces a coloring of  $R_G$ : For every  $v \in V(G)$  and  $n \in \mathbb{N}$  the vertices  $(v, n) \in V(R_G)$  are assigned with the color of  $v$ . Hence  $\chi(R_G) \leq \chi(G)$ .

2. First, observe that  $\text{BBC}(R_G, T_G)$  is finite since  $\text{BBC}(R_G, T_G) \leq 2\chi(R_G) - 1 = 2\chi(G) - 1$ . Let  $k = \text{BBC}(R_G, T_G)$  and let  $\nu$  be a backbone  $k$ -coloring of  $(R_G, T_G)$ . Our goal will be to construct a 2-tuple  $(k+1)$ -coloring  $c$  of  $G$ . First, we define a function  $c': V(G) \rightarrow 2^{[k]} \setminus \{\emptyset\}$ :

$$c'(v) = \{n \in [k] \mid \text{exists } j \in \mathbb{N} : \nu(v, j) = n\}.$$

Now, we define a function  $c: V(G) \rightarrow \binom{[k+1]}{2}$  in the following way:

- $c(v) = \{i, i+1\}$  if  $c'(v) = \{i\}$ .
- $c(v)$  is any 2-element subset of  $c'(v)$  if  $|c'(v)| \geq 2$ .

It remains to show that  $c$  is a 2-tuple coloring of  $G$ . First, observe that  $c'(u) \cap c'(v) = \emptyset$  for every  $uv \in E(G)$ , since  $\{(u, j_1), (v, j_2)\} \in E(R_G)$  for every  $j_1, j_2 \in \mathbb{N}$ . For any  $uv \in E(G)$ , we will show that  $c(u) \cap c(v) = \emptyset$  by considering three cases:

**$|c'(u)| \geq 2$  and  $|c'(v)| \geq 2$ :** Since  $c'(u) \cap c'(v) = \emptyset$ , we infer  $c(u) \cap c(v) = \emptyset$ .

**$c'(u) = \{i\}$  and  $|c'(v)| \geq 2$  (or vice versa):** Since  $c'(u) \cap c'(v) = \emptyset$ , we infer  $i \notin c(v)$ . It remains to show that  $i+1 \notin c(v) \subseteq c'(v)$ . For a contradiction, suppose that  $i+1 \in c(v)$ . Let  $(v, j) \in R_G$  be a vertex such that  $\nu(v, j) = i+1$  and  $(u, j')$  be its neighbor in  $T_G$  due to Lemma 2.2(3). Then  $\nu(u, j') = i$  since  $c'(u) = \{i\}$ . It contradicts the fact that  $\nu$  is a backbone coloring of  $(R_G, T_G)$ .

**$c'(u) = \{i_1\}$  and  $c'(v) = \{i_2\}$ :** Since  $c'(u) \cap c'(v) = \emptyset$ , we have  $i_1 \neq i_2$ . Thus without loss of generality, we can assume that  $i_1 < i_2$ . Moreover,  $i_1 + 1 \notin \{i_2\}$  from a similar reason as in the previous case. Thus,  $i_1 + 1 < i_2$  implies that  $c(u) \cap c(v) = \emptyset$ .

□

### 3 Mycielski-type construction

Mycielski [10] was among the first authors who showed the existence of triangle-free graphs with arbitrarily large chromatic number. We wish, in addition, to




Figure 2: An example of Construction D. In the graph  $G$ , the subgraph  $G[W]$  is indicated by a thick line; circular vertices belong to  $A_1$  and square vertices belong to  $A_2$ .

relate the chromatic number and the 2-tuple chromatic number. More precisely, we will show that for every  $n \in \mathbb{N}$  there exists a triangle-free graph which chromatic number is  $n$  and 2-tuple chromatic is  $2n$ . We will present a construction that increases the chromatic number by 1, the 2-tuple chromatic number by 2, and preserves the property being triangle-free. The construction has several steps.

**Construction D.** Suppose that we are given a graph  $G$ ; a set  $W \subseteq V(G)$  such that  $G[W]$  is bipartite with parts  $A_1$  and  $A_2$  (possibly empty); and two vertices  $z_1, z_2 \notin V(G)$ .

We construct a graph<sup>1</sup>  $D = D(G, W, z_1, z_2)$ . Let<sup>2</sup>  $W' = W \times \{W\}$  be a copy of  $W$  and let  $w'$  be an abbreviation for  $(w, W) \in W'$  where  $w \in W$ . We define

$$\begin{aligned}
 V(D) &= V(G) \cup W' \cup \{z_1, z_2\}, \text{ and} \\
 E(D) &= E(G) \\
 &\cup \{w'_1 w'_2 \mid w_1, w_2 \in W, \text{ and } w_1 w_2 \in E(G)\} \\
 &\cup \{v w' \mid v \in V(G) \setminus W, w \in W, \text{ and } v w \in E(G)\} \\
 &\cup \{w' z_i \mid w \in A_i, i \in \{1, 2\}\}.
 \end{aligned}$$

An example of the construction is depicted on Figure 2. It is easy to check that the following lemma holds:

**Lemma 3.1.** *The graph  $D = D(G, W, z_1, z_2)$  from Construction D has the following properties:*

1. *If  $G$  is triangle-free then  $D$  is triangle-free.*

<sup>1</sup>Formally, the graph  $D$  also depends on a partition of  $W$  to  $A_1$  and  $A_2$ . For our purposes, it will be convenient to suppose that  $W$  is always already given with such a partition.

<sup>2</sup>For the most of the purposes  $W \times \{W\}$  could be replaced by  $W \times \{1\}$ . However it will be convenient later on to get different copies for different  $W$ .






Figure 3: A scheme of Construction H.

2. The graph  $D[(V(G) \setminus W) \cup W']$  is isomorphic to  $G$ .

□

We define another auxiliary graph:

**Construction H.** Suppose that we are given a graph  $G$ , and two vertices  $z_1, z_2 \notin V(G)$ .

We define the graph  $H(G, z_1, z_2)$  in the following way: Order all  $W \subseteq V(G)$  such that  $G[W]$  is bipartite in a sequence  $W_1, W_2, \dots, W_t$  (and choose parts  $A_1, A_2$  for each of them). Construct graphs  $D_i = D(G, W_i, z_1, z_2)$ . Finally, define

$$H = H(G, z_1, z_2) = \bigcup_{i=1}^t D_i,$$

i.e.  $H$  consists of union of sets  $D(G, W_i, z_1, z_2)$  where  $G, z_1$  and  $z_2$  are identified in all the copies, however the sets  $W'_i$  are not identified (see Figure 3).

The following lemma is the key lemma for our construction.

**Lemma 3.2.** *Let  $H = H(G, z_1, z_2)$  be a graph from Construction H.*

1. *If  $G$  is triangle-free, then  $H$  is also triangle-free.*
2. *Let  $k$  be the 2-tuple chromatic number of  $G$ . Then, there is no 2-tuple  $(k+1)$ -coloring  $c$  of  $H$  such that  $c(z_1) = c(z_2)$ .*

*Proof.* The first claim easily follows from Lemma 3.1(1).

For the second claim, assume to the contrary that  $c$  is a 2-tuple  $(k+1)$ -coloring of  $H$  such that  $c(z_1) = c(z_2) = \{k, k+1\}$ . Recall that  $H$  contains  $G$ . Let  $W = \{v \in V(G) \mid c(v) \cap \{k, k+1\} \neq \emptyset\}$ . It is easy to see that  $G[W]$  is a bipartite, thus there exists  $i \in [t]$  (where  $t$  is defined as in Construction H) such




Figure 4: A scheme of Construction J.

that  $W = W_i$ . The graph  $G' = H[(V(G) \setminus W_i) \cup W'_i]$  is isomorphic to  $G$  according Lemma 3.1(2) (where  $W'_i = W_i \times \{W_i\}$  is defined as in Construction D).

We claim that  $c(v) \cap \{k, k+1\} = \emptyset$  for every  $v \in V(G')$ : If  $v \in V(G) \setminus W_i$  then it follows from the definition of  $W = W_i$ , if  $v \in W'_i$  then either  $z_1$  or  $z_2$  is a neighbor of  $v$ . Thus  $c$  restricted to  $G'$  is a 2-tuple  $(k-1)$ -coloring of a graph isomorphic to  $G$  contradicting the assumptions of the lemma.  $\square$

Finally, for a graph  $G$  we define the graph  $J(G)$  that will satisfy our requirements.

**Construction J.** Let  $G$  be a graph,  $k = \chi_2(G)$ ,  $r = \binom{k+1}{2} + 1$ , and let  $Z$  be the graph with  $V(Z) = \{z_1, z_2, \dots, z_r\}$  and  $E(Z) = \emptyset$ . For every  $i, j \in [r]$ ,  $i \neq j$ , let  $G_{ij}$  be an isomorphic copy of  $G$ , formally,  $V(G_{ij}) = V(G) \times \{\{i, j\}\}$  and  $E(G_{ij}) = \{(u, \{i, j\}), v, \{i, j\}\} \mid uv \in E(G)\}$ . Then, we define

$$J(G) = \bigcup_{\{i, j\} \subset [r]} H(G_{ij}, z_i, z_j),$$

i.e.,  $J(G)$  consists of an independent set  $Z$  where between each two vertices  $z_i, z_j$  of  $Z$  there is inserted a copy of  $H(G, z_i, z_j)$ .

**Theorem 3.3.** *Let  $G$  be a graph. The graph  $J(G)$  satisfies the following properties:*

1. *If  $G$  is triangle-free then  $J(G)$  is also triangle-free.*
2.  $\chi_2(J(G)) \geq \chi_2(G) + 2$ .
3.  $\chi(J(G)) \leq \chi(G) + 1$ .

In fact it is not difficult to derive that  $\chi_2(J(G)) = \chi_2(G) + 2$  and  $\chi(J(G)) = \chi(G) + 1$ , but we will not need it for our purposes.

*Proof.* We prove each of the claims separately:

1. This claim follows from Lemma 3.2(1) and from the fact that no two  $z_i$  and  $z_j$  are adjacent in  $H(G_{ij}, z_i, z_j)$ .
2. We use the notation from Construction J. Let  $k = \chi_2(G)$ . We will show that there is no 2-tuple  $(k + 1)$ -coloring  $c$  of  $J(G)$ . For a contradiction, suppose that such  $c$  exists. From the pigeonhole principle, there are  $i, j \in [r]$  such that  $c(z_i) = c(z_j)$ . But it contradicts Lemma 3.2(2) for  $H = H(G_{ij}, z_i, z_j)$ .
3. Again, we use the notation from Construction J. Let  $l = \chi(G)$ , we will show that there is an  $(l + 1)$ -coloring  $\gamma$  of  $J(G)$ . First, we color all the vertices of  $Z$  with color  $l + 1$ , i.e.  $\gamma(Z) = \{l + 1\}$ . Then it is sufficient to color every  $H(G_{ij}, z_i, z_j)$  separately. For notational convenience, we will color  $H(G, z_1, z_2)$  following Construction H so that  $\gamma(z_1) = \gamma(z_2) = l + 1$ . The graph  $G$  is  $l$ -colorable, hence the coloring  $\gamma$  can be extended to  $G$  so that  $\gamma$  is a coloring of  $G$  using only colors  $1, 2, \dots, l$ . Finally, for  $i \in [t]$  and for  $(w, W_i) \in W_i \times \{W_i\}$  we define  $\gamma(w, W_i) = \gamma(w)$ . It is easy to check that  $\gamma$  is an  $(l + 1)$ -coloring of  $H(G, z_1, z_2)$ .

□

**Corollary 3.4.** *For every  $n \in \mathbb{N}$  there exists a (connected) triangle-free graph  $J_n$  such that  $\chi(J_n) = n$  and  $\chi_2(J_n) = 2n$ .*

*Proof.* Let  $J_1$  is the graph consisting of a single vertex. For  $n \geq 2$ , let  $J_n = J^{n-2}(K_2)$ . Theorem 3.3 implies that  $\chi(J_n) \leq n$  and  $\chi_2(J_n) \geq 2n$ , however, it is easy to see that  $\chi_2(G) \leq 2\chi(G)$  for any graph  $G$ . □

The proof of the following corollary, answering negatively Question 1.1, is explicitly written in the introduction (Step III):

**Corollary 3.5.** *For every  $n \in \mathbb{N}$  there exists a (finite) triangle-free graph  $R_n$  and its spanning tree  $T_n$  such that  $\text{BBC}(R_n, T_n) = 2\chi(R_n) - 1 = 2n - 1$ .*

□

## 4 Conclusion

We showed the existence of triangle-free graphs  $R_n$  such that their backbone colorings (with suitable spanning tree) need  $2\chi(R_n) - 1 = 2n - 1$  colors. However, these graphs contain 4-cycles. For further research, it could be interesting to describe the behavior of maximum possible backbone number for graphs with given chromatic number  $\chi$  and given girth  $g$ .

The construction of a graph  $J_n$  can be for every  $t \geq 2$  generalized in an obvious way to get triangle-free graphs  $J_n^t$  such that  $\chi(J_n^t) = n$  and  $\chi_t(J_n^t) = tn$  (compare

with Corollary 3.4). In a bit more detail, to construct graphs  $J_n^t$ , consider  $t$ -colorable subgraphs  $W$  instead of bipartite subgraphs in Constructions D and H and put  $r = \binom{k+t-1}{t} + 1$  in Construction J. Motivated by these results we pose the following conjecture:

**Conjecture.** *For every  $n \in \mathbb{N}$  there exists a finite triangle-free graph  $F_n$  such that  $\chi(F_n) = \chi_f(F_n) = n$ .*

## References

- [1] H. Broersma, F. V. Fomin, P. A. Golovach, and G. J. Woeginger. Backbone colorings for graphs: Tree and path backbones. *J. Graph Theory*, 55(2):137–152, 2007.
- [2] H. J. Broersma, J. Fujisava, L. Marchal, D. Paulusma, A. N. M. Salman, and K. Yoshimoto.  $\lambda$ -backbone colorings along pairwise disjoint stars and matchings. Manuscript, 2007.
- [3] H. J. Broersma, J. Fujisawa, and K. Yoshimoto. Backbone colorings along perfect matchings. Memorandum 1706, Enschede, 2003.
- [4] H. J. Broersma, L. Marchal, and D. Paulusma. Backbone colorings along stars and matchings in split graphs: their span is close to the chromatic number. Manuscript, 2007.
- [5] P. Erdős. Graph theory and probability. *Canad. J. Math.*, 11:34–38, 1959.
- [6] P. Hell and J. Nešetřil. *Graphs and homomorphisms*. Oxford University Press, 2004.
- [7] M. Larsen, J. Propp, and D. Ullman. The fractional chromatic number of Mycielski’s graphs. *J. Graph Theory*, 19:411–416, 1995.
- [8] J. Matoušek. *Using the Borsuk-Ulam Theorem*. Springer, Berlin etc., 2003.
- [9] J. Miškuf, R. Škrekovski, and M. Tancer. Backbone colorings of graphs with bounded maximum degree. Manuscript, 2008.
- [10] J. Mycielski. Sur le coloriage des graphes. *Colloq. Math.*, 3:161–162, 1955.