

UNIVERSITY OF LJUBLJANA  
INSTITUTE OF MATHEMATICS, PHYSICS AND MECHANICS  
DEPARTMENT OF MATHEMATICS  
JADRANSKA 19, 1000 LJUBLJANA, SLOVENIA

**Preprint series, Vol. 46 (2008), 1046**

MEDIAN GRAPHS, THE  
REMOTENESS FUNCTION,  
PERIPHERY TRANSVERSALS,  
AND GEODETIC NUMBER TWO

Kannan Balakrishnan    Boštjan Brešar  
Manoj Changat        Wilfried Imrich  
Sandi Klavžar        Matjaž Kovše  
Ajitha R. Subhamathi

ISSN 1318-4865

March 25, 2008

Ljubljana, March 25, 2008

# Median graphs, the remoteness function, periphery transversals, and geodetic number two\*

Kannan Balakrishnan<sup>†</sup>    Boštjan Brešar<sup>‡</sup>    Manoj Changat<sup>§</sup>  
Wilfried Imrich<sup>¶</sup>    Sandi Klavžar<sup>||</sup>    Matjaž Kovše<sup>\*\*</sup>  
Ajitha R. Subhamathi<sup>††</sup>

## Abstract

A periphery transversal of a median graph  $G$  is introduced as a set of vertices that meets all the peripheral subgraphs of  $G$ . Using this concept, median graphs with geodetic number 2 are characterized in two ways. They are precisely the median graphs that contain a periphery transversal of order 2 as well as the median graphs for which there exists a profile such that the remoteness function is constant on  $G$ . Moreover, an algorithm is presented that decides in  $O(m \log n)$  time whether a given graph  $G$  with  $n$  vertices and  $m$  edges is a median graph with geodetic number 2. Several additional structural properties of the remoteness function on hypercubes and median graphs are obtained and some problems listed.

**Keywords:** median graph; median set; remoteness function; geodetic number; periphery transversal; hypercube

**AMS Subj. Class. (2000):** Primary: 05C85; Secondary: 05C12, 90B80;

---

\*Work supported by the Ministry of Science of Slovenia and by the Ministry of Science and Technology of India under the bilateral India-Slovenia grants BI-IN/06-07-002 and DST/INT/SLOV-P-03/05, respectively.

<sup>†</sup>Department of Computer Applications, Cochin University of Science and Technology, Cochin-22, India, bkannan@cusat.ac.in

<sup>‡</sup>University of Maribor, FEECS, Smetanova 17, 2000 Maribor, Slovenia, bostjan.bresar@uni-mb.si

<sup>§</sup>Department of Futures Studies, University of Kerala, Trivandrum-695034, India, mchangat@gmail.com

<sup>¶</sup>Department of Mathematics and Information Technology, Montanuniversität Leoben, Austria, wilfried.imrich@mu-leoben.at

<sup>||</sup>Department of Mathematics and Computer Science, FNM, University of Maribor, Gosposvetska 84, 2000 Maribor, Slovenia, sandi.klavzar@uni-mb.si

<sup>\*\*</sup>Institute of Mathematics, Physics and Mechanics, Jadranska 19, 1000 Ljubljana, Slovenia, matjaz.kovse@gmail.com

<sup>††</sup>Department of Futures Studies, University of Kerala, Trivandrum-695034, India, ajithars@gmail.com

# 1 Introduction

Given a profile (that is, a multiset of vertices) on a graph, the location theory quests for vertices whose remoteness (the sum of distances to the vertices of the profile) is minimum or maximum, and these sets are called median and antimedian sets, respectively. The problem of locating median sets for profiles on graphs was considered by many authors; see, for example, [2, 4, 5, 15, 16]. On the other hand, not much work has been done so far on the antimedian problem for profiles on graphs, and though the two problems look similar, there are important differences. For instance, while it is clear that any vertex can be in the median set of a graph for some profile, this is not always true for the antimedian set.

In this paper we give a closer look at the remoteness function in median graphs with the aim to shed more light on the antimedian problem in this class. Median graphs form a closely investigated and well understood class of graphs, and are probably the most important class of graphs in metric graph theory (we refer to a comprehensive survey on median graphs [13]). Hence it is not surprising that they were investigated also in location theory. For instance, it is known that in median graphs median sets are always intervals between two vertices [4], and in particular, for any odd profile they consist of exactly one vertex [15].

We show in this paper that for any odd profile the antimedian set is an independent set of vertices that lie in a strict boundary of a median graph. On the other hand, it can happen in a special class of median graphs that the entire vertex set is the antimedian set of some even profile. These are precisely the median graphs with geodetic number 2, that were studied previously in [6], where several characterizations of these graphs were obtained. In this paper we add two more characterizations, and one of them is used in the algorithm for the recognition of median graphs with geodetic number 2 in Section 4.

In the next section we fix the notation and state some preliminary results. In Section 3 we prove two characterizations of median graphs with geodetic number two, one of which involves the remoteness function. In addition we obtain some properties of antimedian sets in median graphs. Section 4 is concerned with an algorithm for the recognition of median graphs with geodetic number 2. Median graphs are a subclass of the class of isometric subgraphs of hypercubes. The complexity of recognizing whether a given graph  $G$  with  $n$  vertices and  $m$  edges is such a graph is  $O(mn)$  in general. For median graphs this essentially reduces to  $O(m\sqrt{n})$ ; see [9]. There is little hope to reduce it further in general, since it is closely related to that of recognizing triangle-free graphs (see [12]). However, in special cases the complexity is much lower. For example, it is  $O(m)$  for planar median graphs. Here we show that median graphs with geodetic number 2 can be recognized in  $O(m \log n)$  time.

In Section 5 we study the remoteness function in hypercubes (and Hamming graphs) which is then used in the final section. There a theorem is proved which establishes a connection between antimedian sets on a median graph  $G$  and antimedian

sets on the hypercube, into which  $G$  is embedded isometrically.

## 2 Preliminaries

In this paper we consider simple, undirected, finite, and connected graphs. The *distance* considered in this paper is the usual shortest path distance  $d$ . A shortest path between vertices  $u$  and  $v$  will be called a  $u, v$ -*geodesic*. The set of vertices on all  $u, v$ -geodesics is called the *interval* between  $u$  and  $v$ , denoted  $I(u, v)$ . A set  $S$  of vertices in a graph  $G$  is called the *geodetic set* of  $G$  if for every vertex  $x \in V(G)$  there exist  $u, v \in S$  such that  $x \in I(u, v)$ . The *geodetic number*  $g(G)$  of a graph  $G$  is the least size of a set of vertices  $S$  such that any vertex from  $G$  lies on a  $u, v$ -geodesic, where  $u, v \in S$ . For a connected graph  $G$  and subsets of vertices  $X, Y \subseteq V(G)$  we will write  $d(X, Y) = \min\{d(x, y) \mid x \in X, y \in Y\}$ . In particular, for a vertex  $u$  of  $G$  and a set of vertices  $X$  we have  $d(u, X) = \min\{d(u, x) \mid x \in X\}$ .

A *profile*  $\pi = (x_1, \dots, x_k)$  on a graph  $G$  is a finite sequence of vertices of  $G$ , and  $k = |\pi|$  is called the *size* of the profile  $\pi$ . Note that in a profile a vertex may be repeated. Given a profile  $\pi$  on  $G$  and a vertex  $u$  of  $G$ , the *remoteness*  $D(u, \pi)$  (see [14]) is

$$D(u, \pi) = \sum_{x \in \pi} d(u, x).$$

The vertex  $u$  is called a *median (antimedian) vertex* for  $\pi$  if  $D(u, \pi)$  is minimum (maximum). The *median (antimedian) set*  $M(\pi, G)$  ( $AM(\pi, G)$ ) of  $\pi$  in  $G$  is the set of all median (antimedian) vertices for  $\pi$ .

A (connected) graph  $G$  is a *median graph* if for any three vertices  $x, y, z$  there exists a unique vertex that lies on geodesics between all pairs of  $x, y, z$ . Two of the most important classes of median graphs are trees and hypercubes. The *hypercube* or  $n$ -*cube*  $Q_n$ ,  $n \geq 1$ , is the graph with vertex set  $\{0, 1\}^n$ , two vertices being adjacent if the corresponding tuples differ in precisely one position. A vertex  $u$  of  $Q_n$  will be written in its coordinate's form as  $u = u^{(1)} \dots u^{(n)}$ . A natural generalization of hypercubes are *Hamming graphs*, whose vertices are  $m$ -tuples  $u = u^{(1)} \dots u^{(m)}$ , such that  $0 \leq u^{(i)} \leq m_i - 1$ , where  $m_i \geq 2$  for each  $i$ , and adjacency is defined in the same way (that is, two vertices are adjacent precisely when they differ in exactly one coordinate). Note that the distance between vertices in Hamming graphs coincides with the Hamming distance (that is, the number of coordinates in which the  $m$ -tuples differ).

A subgraph  $H$  of a (connected) graph  $G$  is an *isometric subgraph* if  $d_H(u, v) = d_G(u, v)$  holds for any vertices  $u, v \in H$ . Let  $G$  be an isometric subgraph of some hypercube. The smallest integer  $d$  such that  $G$  is an isometric subgraph of  $Q_d$  is called the *isometric dimension* of  $G$  and denoted  $\text{idim}(G)$ . An important structural result due to Mulder [17] asserts that every median graph  $G$  can be isometrically embedded in a hypercube such that the median of every profile  $\pi$  of cardinality three

in  $G$  on the hypercube coincides with the median of  $\pi$  in  $G$ . A subset  $S$  of vertices in a graph  $G$  is *convex* in  $G$  if  $I(u, v) \subseteq S$  for any  $u, v \in S$ . It is well-known that convex sets in median graphs enjoy the *Helly property*, that is, any family of pairwise disjoint convex sets has a common intersection.

For a connected graph and an edge  $xy$  of  $G$  we denote

$$W_{xy} = \{w \in V(G) \mid d(x, w) < d(y, w)\}.$$

Note that if  $G$  is a bipartite graph then  $V(G) = W_{ab} \cup W_{ba}$  holds for any edge  $ab$ . Next, for an edge  $xy$  of  $G$  let  $U_{xy}$  denote the set of vertices  $u$  that are in  $W_{xy}$  and have a neighbor in  $W_{yx}$ . Sets in a graph that are  $U_{xy}$  for some edge  $xy$  will be called *U-sets*. Similarly we define *W-sets*. If for some edge  $xy$ ,  $W_{xy} = U_{xy}$ , we call the set  $U_{xy}$  *peripheral set* or *periphery*.

Edges  $e = xy$  and  $f = uv$  of a graph  $G$  are in the Djoković-Winkler relation  $\Theta$  [8, 21] if  $d_G(x, u) + d_G(y, v) \neq d_G(x, v) + d_G(y, u)$ . Relation  $\Theta$  is reflexive and symmetric. If  $G$  is bipartite, then  $\Theta$  can be defined as follows:  $e = xy$  and  $f = uv$  are in relation  $\Theta$  if  $d(x, u) = d(y, v)$  and  $d(x, v) = d(y, u)$ . It is well-known that the relation  $\Theta$  is transitive in isometric subgraphs of hypercubes, and so it is an equivalence relation on the edge set of every median graph. Note that peripheral sets are precisely the  $U$ -sets that induce a connected component of  $G - F$  for some  $\Theta$ -class  $F$ .

### 3 Median graphs with geodetic number two

In this section we characterize median graphs with geodetic number 2. One characterization is described in terms of so-called periphery transversal, a concept that could be of independent interest in the study of median graphs. The other characterization answers the following question from location theory: for which median graphs  $G$  their vertex-set is the (anti)median set of some profile on  $G$ .

Let  $G$  be a median graph. We say that a set  $S$  is a *periphery transversal* if every peripheral subgraph of  $G$  contains a vertex of  $S$ . It was proved in [6] that every geodetic set is periphery transversal set. Let  $\text{pt}(G)$  denote the size of a minimum periphery transversal in a median graph  $G$ . Then, clearly,  $\text{pt}(G) \leq g(G)$  for any median graph  $G$ . On the other hand, it may happen that any minimum geodetic set of a median graph  $G$  must contain some vertices that are not in a peripheral subgraph. For instance, in the graph  $G$  obtained from the 3-cube by attaching a leaf to 3 independent vertices we have  $\text{pt}(G) = 3 < 4 = g(G)$ .

We need the following well-known facts, see [11].

**Lemma 3.1** *Let  $G$  be a median graph,  $C$  a cycle,  $P$  a geodesic, and  $F$  a  $\Theta$ -class of  $G$ . Then*

$$(i) \quad F \cap C \neq \emptyset \Rightarrow |F \cap C| \geq 2;$$

(ii)  $F \cap P \neq \emptyset \Rightarrow |F \cap P| = 1$ .

We also recall the following theorem from [6].

**Theorem 3.2** *Let  $G$  be a median graph. Then  $g(G) = 2$  if and only if there exist vertices  $a, b \in V(G)$  and an  $a, b$ -geodesic that contains edges from all  $\Theta$ -classes of  $G$ .*

Combining Lemma 3.1 with Theorem 3.2 we infer that if  $a$  and  $b$  are as in the theorem above, then on any geodesic from  $a$  to  $b$  all  $\Theta$ -classes appear. Conversely,  $g(G) > 2$  implies that for any two vertices  $a$  and  $b$  in  $G$  there exists a  $\Theta$ -class whose edges are outside  $I(a, b)$ .

**Theorem 3.3** *For a median graph  $G$  the following statements are equivalent.*

- (i)  $g(G) = 2$ ,
- (ii)  $\text{pt}(G) = 2$ ,
- (iii)  $D(x, \pi)$  is constant on  $G$  for some profile  $\pi$ .

**Proof.** (i) $\Rightarrow$ (ii): Let  $G$  be a median graph with  $g(G) = 2$ . As every  $W$ -set in a median graph contains a periphery, we infer that  $\text{pt}(G) \geq 2$ . We have already observed that in general  $\text{pt}(G) \leq g(G)$ , hence  $\text{pt}(G) = 2$ .

(ii) $\Rightarrow$ (i): Let  $G$  be a median graph with  $\text{pt}(G) = 2$ , and assume to the contrary that  $g(G) > 2$ . Then for any two vertices  $a, b \in V(G)$ ,  $I(a, b) \neq V(G)$ , and by Theorem 3.2 we infer that there exists a  $\Theta$ -class  $F$  that lies outside  $I(a, b)$ . Then there also exists a  $W$ -set  $W_{xy}$  that has an empty intersection with  $I(a, b)$ . In addition,  $W_{xy}$  contains a periphery that does not contain  $a$  and  $b$ . Thus  $\{a, b\}$  is not a periphery transversal, and since  $a$  and  $b$  were chosen arbitrarily we infer that  $\text{pt}(G) > 2$ , a contradiction.

(i) $\Rightarrow$ (iii): Let  $a$  and  $b$  be vertices in  $G$  such that  $I(a, b) = V(G)$ . Set  $\pi = (a, b)$ . Since for any  $x \in V(G)$  we have  $d(a, x) + d(x, b) = d(a, b) = \text{diam}(G)$  we get  $D(x, \pi) = \text{diam}(G)$ .

(iii) $\Rightarrow$ (i): For this direction we recall a result by Bandelt and Barthélemy [4, Proposition 6] which says that for any profile  $\pi$  on a median graph  $G$ , the median set  $M(\pi, G)$  coincides with the interval  $I(\alpha(\pi), \beta(\pi))$  (where  $\alpha(\pi)$  and  $\beta(\pi)$  are two vertices in  $G$  obtained by a formula in the associated median semilattice). Hence, if  $D(x, \pi)$  is constant on  $G$  for a profile  $\pi$ , then  $V(G) = M(\pi, G) = I(\alpha(\pi), \beta(\pi))$ , which in turn implies  $g(G) = 2$ .  $\square$

By the above theorem, in a median graph  $G$  the whole vertex set is (anti)median if and only if  $g(G) = 2$ . However, even if  $g(G) > 2$ , every vertex can be in some median set of  $G$  (e.g., by taking this vertex as the unique vertex in the profile). We believe that the antimedian case is different, and suspect that if a median graph  $G$

has geodetic number greater than two then there are vertices that cannot lie in the antimedian set for any profile on  $G$ . We present two partial results that confirm this; in the first one we show this for an arbitrary odd profile.

A vertex  $v$  of a graph  $G$  is a *strict boundary vertex* (with respect to  $v'$ ) of  $G$  if there exists a vertex  $v'$  such that for any neighbor  $u$  of  $v$ ,  $d(v', v) > d(v', u)$  (in other words, the neighborhood of  $v$  is contained in  $I(v, v')$ ). The *strict boundary*  $\bar{\partial}G$  of a graph  $G$  is the set of strict boundary vertices in  $G$ .

For an edge  $uv$  in a median graph  $G$  and a profile  $\pi$ , we let  $\pi_{uv} = W_{uv} \cap \pi$ . As usually,  $|\pi_{uv}|$  denotes the size of the profile  $\pi$  in  $W_{uv}$ . Note that  $|\pi_{uv}| > |\pi_{vu}|$  implies that the median set of  $\pi$  on  $G$  lies in  $W_{uv}$  which in turn implies that if  $u$  and  $v$  are both in a median set then  $|\pi_{uv}| = |\pi_{vu}|$ . These observations are a basis for several strategies to find median sets in median-like graphs, see [2, 16].

**Lemma 3.4** *Let  $\pi$  be an odd profile in a median graph  $G$ , then every vertex in  $AM(\pi, G)$  is a strict boundary vertex.*

**Proof.** Let  $v \in AM(\pi, G)$  and  $|\pi|$  be odd. Since for every neighbor  $u_i$  of  $v$ ,  $|\pi_{u_i v}| > |\pi_{v u_i}|$  we infer that

$$|\pi_{u_i v}| > \frac{|\pi|}{2}.$$

Hence  $\pi_{u_i v}$  and  $\pi_{u_j v}$  intersect for any neighbors  $u_i, u_j$  of  $v$ , where  $i \neq j$ . Since  $\pi_{u_i v} \subseteq W_{u_i v}$ , the sets  $W_{u_i v}$  also pairwise intersect for all neighbors  $u_i$  of  $v$ . Since  $W$ -sets are convex, by the Helly property for convex sets there exists a vertex

$$v' \in \bigcap_{u_i \in N(v)} W_{u_i v}.$$

Hence  $u_i$  is strictly closer to  $v'$  than  $v$  for any  $i$ , and so  $v$  is a strict boundary vertex (with respect to  $v'$ ).  $\square$

From the proof of the lemma above we also see that no neighbor of  $v \in AM(\pi, G)$  achieves  $D(v, \pi)$ , hence we derive the following result.

**Proposition 3.5** *Let  $\pi$  be an odd profile in a median graph  $G$ . Then  $AM(\pi, G)$  is an independent set in  $G$  and  $AM(\pi, G) \subseteq \bar{\partial}G$ .*

We leave the structure of antimedian sets for even profiles in median graphs as an open problem. Note that in this case antimedian vertices need not be in a strict boundary, even if  $g(G) > 2$ . For instance, let  $G$  be obtained from the  $3 \times 3$  grid (that is the Cartesian product  $P_3 \square P_3$ ) so that to the central vertex another vertex  $a$  is attached, and let the profile  $\pi$  consist of two vertices  $u, v$  of degree two such that  $d(u, v) = 2$ . Then  $AM(\pi, G) = \{x, y, z, a\}$ , where  $x$  and  $y$  are another two vertices of degree two (different from  $u$  and  $v$ ), and  $z$  is their common neighbor. Note that  $z$  is not a strict boundary vertex in  $G$ . However, we suspect that the following question has affirmative answer.

**Question 3.6** *Let  $G$  be a median graph and  $g(G) > 2$ . Is it true that there exists a vertex in  $G$  that is not in  $AM(\pi, G)$  for all profiles  $\pi$  on  $G$ ?*

We end this section with another result that partially confirms the positive answer to the above question. It describes the antimedian set in median graphs in the special case when the profile is the whole vertex set, each vertex appearing exactly once. This problem is known in the literature as the obnoxious center problem, and has been quite well studied, cf. [7, 19, 20, 22].




Figure 1: An antimedian vertex that is not peripheral.

In the case when the profile is the vertex set, one might ask the following question: is it true that every antimedian vertex lies in a periphery of a median graph? The answer is negative, as can be seen in the example from Fig. 1. The black vertex is the unique antimedian vertex of this graph, as soon as there are sufficiently many gray pendant vertices. Note that the black vertex is not in any periphery of this median graph. However, we can prove a result similar to Proposition 3.5.

**Proposition 3.7** *Let  $G$  be a median graph, and let  $\pi$  be the profile, consisting of vertices of  $V(G)$  (with no repetitions). If  $v \in AM(\pi, G)$  then  $v$  is a strict boundary vertex.*

**Proof.** Let  $v \in AM(\pi, G)$ . We infer that for every neighbor  $u_i$  of  $v$ ,  $|W_{u_i v}| \geq |W_{v u_i}|$ , hence

$$|W_{u_i v}| \geq \frac{|V(G)|}{2}.$$

Let  $u_1, \dots, u_t$  be the neighbors of  $v$ . If  $t = 1$ , that is,  $v$  has only one neighbor, then  $v$  is clearly a strict boundary vertex with respect to any other vertex. Suppose that

$u_i, u_j$  are neighbors of  $v$  and  $i \neq j$ . Then by the above

$$|W_{u_i v}| + |W_{u_j v}| \geq |V(G)|.$$

Since  $v \notin W_{u_i v}$ , for any  $i$ , we find that  $W_{u_i v}$  and  $W_{u_j v}$  intersect. Since  $W$ -sets are convex, we infer by the Helly property for convex sets that there exists a vertex

$$v' \in \bigcap_{i=1}^t W_{u_i v}.$$

Hence  $v$  is a strict boundary vertex with respect to  $v'$  which completes the proof of the proposition.  $\square$

## 4 Recognition of median graphs with geodetic number two

As already mentioned, median graphs are isometric subgraphs of hypercubes (*partial cubes* for short), and the recognition complexity for such graphs is  $O(mn)$ . In other words, there exists an algorithm that recognizes whether any given graph  $G$  with  $n$  vertices and  $m$  edges is a partial cube in  $O(mn)$  time. The algorithm also provides an embedding of  $G$ . In the rest of this section  $n$  and  $m$  will denote the number of vertices and edges of a given graph.

However, if it is known that a graph  $G$  is a median graph, then  $G$  can be embedded isometrically into a hypercube in  $O(m \log n)$  time. This discrepancy between the embedding complexity and the recognition complexity was a strong motivation to find better recognition algorithms for median graphs. The algorithm of Hagauer, Imrich and Klavžar [9] with complexity  $O(m\sqrt{n})$  was the first of this kind. Later Imrich [11, Theorem 7.27] derived the asymptotically better result  $O((m \log n)^{1.41})$ . Here the exponent 1.41 actually is  $2\omega/(\omega+1)$ , where  $\omega$  is the exponent of matrix multiplication with its current value 2.376. By a result of Imrich, Klavžar and Mulder [12] this recognition complexity is closely related with the recognition complexity of triangle-free graphs. Hence improvements of the recognition complexity of median graphs seem to be very difficult.

Nonetheless, some classes of median graphs can be recognized much faster. This includes planar median graphs [12], which can be recognized in linear time and acyclic cubical complexes [10], which can be recognized in  $O(m \log n)$  time. Here we show that median graphs with geodetic number two can also be recognized in  $O(m \log n)$  time. This is possible because of a bound on the maximum degree of a median graph with geodetic number two and the fact that every peripheral subgraph meets geodetic set, see Brešar and Tepoh Horvat [6].

We begin with the bound on the maximum degree  $\Delta(G)$  of a median graph  $G$  with  $g(G) = 2$ .

**Lemma 4.1** *Let  $G$  be a median graph with  $g(G) = 2$ . Then  $\Delta(G) \leq 2 \log_2 n$ .*

**Proof.** Suppose  $G = I_G(v, w)$  and let  $L_0, L_1, \dots, L_r$  be the levels of the BFS-ordering of the vertices of  $G$  with respect to a root  $v$ ; see e.g. [11, p. 41]. Let  $x \in L_i$  and  $xy \in E(G)$ . Since  $G$  is bipartite  $y \notin L_i$ . If  $y \in L_{i-1}$  we call the edge  $xy$  a *down-edge* and otherwise an *up-edge*. Clearly  $y$  is closer to  $v$  than  $x$  if  $xy$  is a down-edge, and closer to  $w$  if  $xy$  is an up-edge. In other words, the up-edges with respect to  $v$  are the down-edges with respect to  $w$ . By [11, Lemma 3.35] the number of down-edges of every vertex  $x$  in a median graph is bounded by  $\log_2 n$ . Clearly the number of up-edges satisfies the same bound, hence  $d(v) \leq 2 \log_2 n$  for all  $v \in V(G)$ .  $\square$

Next we show how to check efficiently whether a given induced subgraph of a graph  $G$  is also a convex subgraph. For a subgraph  $H$  of a graph  $G$  let  $\partial H$  be the set of edges with one endvertex in  $H$  and the other in  $G \setminus H$ .

**Lemma 4.2** *Let  $H$  be an induced connected subgraph of a partial cube for which the  $\Theta$ -classes are already known. Then the complexity of recognizing whether  $H$  is a convex subgraph of  $G$  is  $O(|E(H)| + |\partial H|)$ .*

**Proof.** By the convexity lemma [11, Lemma 2.7] it suffices to show that no edge of  $\partial H$  is in the relation  $\Theta$  with an edge of  $H$ . In other words, we have to show that the list of  $\Theta$ -classes that meet  $E(H)$  is disjoint from the list of  $\Theta$ -classes that meet  $\partial H$ .

Let  $E_1, \dots, E_k$ , where  $k < n$ , be the  $\Theta$ -classes of  $G$  and  $\mathbf{v}_H$  the 0,1-vector of length  $k$  with  $\mathbf{v}_H(i) = 0$  if  $E_i \cap E(H) = \emptyset$  and  $\mathbf{v}_H(i) = 1$  otherwise. Since the  $\Theta$ -classes are known, we can assume that there exists a function  $c : E(G) \rightarrow \{1, \dots, k\}$  that computes the index  $i$  for which  $e \in E_i$  in constant time. With a well known trick, see [1], the vector  $\mathbf{v}_H$  can be determined in  $O(|E(H)|)$  time, even if  $|E(H)| < k$ , by scanning all edges of  $H$ . Moreover we scan all edges of  $\partial H$ . If  $e \in E_i$  and  $\mathbf{v}_H(i) = 1$ , then  $H$  is not convex. We thus have to check whether  $\mathbf{v}_H(c(e)) = 0$  for all  $e \in \partial H$ . Clearly this can be done in  $O(|\partial H|)$  time.  $\square$

Next we show how to efficiently check the convexity of  $U$ -sets.

**Corollary 4.3** *Let  $H$  be a partial cube for which the  $\Theta$ -classes are already known, and  $\Delta$  the maximum degree of vertices in  $G$ . Then one can check in  $O(m\Delta + m \log n)$  time whether all  $U$ -sets are convex.*

**Proof.** First note that the total size of  $U$ -sets in  $G$  is  $m$ . Furthermore  $|E(U_{ab})| < |U_{ab}| \log_2 |U_{ab}|$  by Graham's density lemma [11, Proposition 1.24]. Hence, for the total number of edges in the  $U$ -sets we have the following inequality

$$\left(\sum |U_{ab}|\right) \max(\log_2 |U_{ab}|) \leq m \log_2 n.$$

Let  $\mathbf{v}_{U_{ab}}$  be defined as in Lemma 4.2. Then it is clear that the set of vectors  $\mathbf{v}_{U_{ab}}$  can be determined in  $O(m \log n)$  time. Since the total size of the sets  $\partial U$  over all  $U$ -sets is bounded by  $m\Delta$  the corollary follows.  $\square$

**Proposition 4.4** *Let  $G$  be a graph with  $\Delta(G) \leq 2 \log_2 n$ . Then one can check in  $O(m \log n)$  time whether  $G$  is a median graph, determine all  $\Theta$ -classes and all  $U$ -sets.*

**Proof.** By [11, Lemma 7.15] one can check in  $O(m \log n)$  time whether  $G$  is a partial cube, determine all  $\Theta$ -classes and all  $U$ -sets. By [11, Corollary 2.27] a partial cube is a median graph if and only if all  $U$ -sets are convex. Now the proof is completed by the observation that the convexity of the  $U$ -sets of a given partial cube can be checked in  $O(m \log n)$  by Corollary 4.3.  $\square$

Next we describe a procedure which can be used to construct all median graphs. For a connected graph  $H$  and its convex subgraph  $P$  the *peripheral expansion of  $H$  along  $P$*  is the graph  $G$  obtained as follows. Let  $P'$  be an isomorphic copy of  $P$  and  $\alpha$  a corresponding isomorphism. Take the disjoint union  $H + P'$  and join each vertex  $v \in P$  by an edge with  $\alpha(v) \in P'$ . We call the new graph a *peripheral expansion of  $H$  along  $P$*  and denote it by  $G = pe(H; P)$ . Mulder [18] proved that a graph is a median graph if and only if it can be obtained from  $K_1$  by a sequence of peripheral expansions.

We still have to find a geodetic set consisting of two elements. In order to accomplish this, we will use this sequence of peripheral expansions to determine all geodetic sets. We begin with a relationship between the geodetic sets of a median graph  $H$  and the graph  $G = pe(H, P)$ .

**Lemma 4.5** *Let  $G = pe(H; P)$  be a median graph and  $\{x, y\}$  a geodetic set of  $H$ , where  $y \in P$ . Then the set  $\{x, z\}$ , where  $z$  is the neighbor of  $y$  in  $G \setminus H$  is a geodetic set in  $G$ . Moreover, all minimum geodetic sets of  $G$  are of this form.*

**Proof.** We have to show that every vertex  $w$  of  $G$  is on a shortest  $xz$ -path. Suppose first  $w \in H$ . Then, clearly  $w$  is on a  $xy$ -geodesic, since  $\{x, y\}$  is a geodetic set in  $H$ . Thus  $w$  is also on  $xz$ -geodesic going through  $y$ . Suppose next  $w \in G \setminus H$  and let  $w'$  be a neighbor of  $w$ , where  $w' \in H$ . Then  $w'$  lies on  $xy$ -geodesic. Let  $L_1$  denote the  $yw'$ -geodesic and let  $L_2$  denote the  $w'x$ -geodesic. Since  $P$  is a convex subgraph of  $H$  (and therefore also of  $G$ )  $L_1$  is completely contained in  $P$ . Recall that in median graph for any edge  $ab$  we have  $U_{ab} \cong U_{ba}$  and that the isomorphism is induced by the edges between  $U_{ab}$  and  $U_{ba}$ . Let  $L'_1$  be the projection of  $L_1$  into  $P'$  by this isomorphism. Then  $L'_1 \cup ww' \cup L_2$  is a  $xz$ -geodesic in  $G$  containing  $w$ . Conversely if  $\{x, z\}$  is a geodetic set in  $G = pe(H; P)$  then by [6, Lemma 2]  $x$  or  $z$

must be in  $P'$ . Suppose  $z$  is in  $P'$ . Then we can use the same arguments as above to see that  $\{x, y\}$  is a geodetic set in  $H$ , where  $y$  is a neighbor of  $z$  in  $H$ .  $\square$

If  $\{x, y\}$  is a geodetic set in  $G$  then this is the only minimum geodetic set containing  $x$ , since by Lemma 4.5  $x$  is uniquely determined by  $y$  and vice versa.

**Corollary 4.6** *Let  $G = pe(H; P)$  be a median graph with  $g(G) = 2$ . Then all minimum geodetic sets of  $G$  can be obtained from the minimum geodetic sets of  $H$  in  $O(|P|)$  time.*

**Proof.** Let  $P' = U_{ab}$ , where  $a \in G \setminus H$ . To find the geodetic sets of  $G$  we scan all vertices  $z$  of  $U_{ab}$ . If the neighbor  $y$  of  $z$  in  $U_{ba}$  is in the geodetic set  $\{y, x\}$  of  $H$ , then by Lemma 4.5  $\{z, x\}$  is a geodetic set of  $G$ . Clearly the complexity of this task is  $O(|U_{ab}|)$ .  $\square$

**Corollary 4.7** *Let  $G$  be a median graph with  $g(G) = 2$ . If the representation of  $G$  as a series of peripheral expansions, starting from  $K_1$ , is known, then all minimum geodetic sets of  $G$  can be obtained in  $O(n)$  time.*

**Proof.** At every expansion step  $|U_{ab}|$  vertices are added at a total cost of  $O(|U_{ab}|)$ . The observation that  $n - 1$  vertices are added altogether completes the proof.  $\square$

We are thus left with the task of representing  $G$  by a series of peripheral expansions.

**Theorem 4.8** *Let  $G$  be a median graph with  $\Delta(G) \leq 2 \log_2 n$ . Then a representation of  $G$  by a series of peripheral expansions can be found in  $O(m \log n)$  time.*

**Proof.** By [11, Lemma 7.15] and Proposition 4.4 we know that one can recognize  $G$  as a median graph, partition its edge set into  $\Theta$ -classes, and determine all  $U$ -sets in  $O(m \log n)$  time. We show now that we can determine all peripheral  $U$ -sets within the same time complexity. We first observe that the peripheral  $U$ -sets are characterized by the fact that  $\partial U$  consist of  $|U|$  independent edges that meet every vertex of a  $U$ -set. In other words  $U_{ab}$  is peripheral if

$$\deg_G(v) = \deg_{U_{ab}}(v) + 1,$$

for every  $v \in U_{ab}$ . Clearly  $\deg_{U_{ab}}(v) + 1 \leq \deg_G(v)$  for  $v \in G$ . Thus, setting

$$ex_{U_{ab}}(v) = \deg_G(v) - \deg_{U_{ab}}(v) - 1$$

it is clear that  $U_{ab}$  is peripheral if and only if

$$ex(U_{ab}) = \sum_{v \in U_{ab}} ex_{U_{ab}}(v) = 0.$$

Intuitively,  $ex(v)$  is the *excess* of the degree of  $v$  above its minimum.

We thus need the degrees of every vertex in its  $U$ -sets and in  $G$ . The degrees of all vertices from a given  $U$ -set  $U_{xy}$  can be determined in  $|E(U_{xy})|$  time and the degrees of all vertices in  $G$  in  $O(m)$  time. Since the total number of edges in the  $U$ -sets is  $O(m \log n)$  we can thus determine all degrees in  $O(m \log n)$  time.

In a second run, scanning all vertices in the  $U$ -sets, we determine excesses of all vertices of  $G$  and calculate the sum of all corresponding excesses of vertices from some  $U$ -set. Since the total number of vertices in the  $U$ -sets is  $O(m)$ , this can be done in the required time too.

In this process we keep a record of all these numbers and consider the first peripheral set we find, say  $U_{ab}$ .

We now show that we can remove  $U_{ab}$  from  $G$  and determine for  $H = G \setminus U_{ab}$  the same data structure we had for  $G$ . In other words, we can determine the adjacency list of all new  $U$ -sets in the graph  $H$ , all degrees and the new values of the excess numbers for all vertices in  $H$  and all the new  $U$ -sets in  $O(|U_{ab}| \log n)$  time.

We first find the new adjacency list of the new  $U$ -sets of  $H$ . We first recall that the removal of a vertex  $v$  and all incident edges from a graph is of complexity  $O(\deg(v))$  if the graph is represented by an extended adjacency list or the adjacency matrix; see pp. 37 in [11]. In  $G$  every vertex  $v$  is also a vertex of every  $U_{vw}$ , where  $w$  is a neighbor of  $v$  in  $G$ . Thus every  $v \in U_{ab}$  is in at most  $O(\log n)$  sets  $U_{vw}$ . The degree of the vertex  $v$  in such a  $U_{vw}$  is  $\deg_G(v) - 1 = \deg_{U_{ab}}(v)$ . The cost of removing  $v$  from all  $U_{vw}$  is thus  $O(\deg_{U_{ab}}(v) \log n)$ . For all  $v \in U_{ab}$  this amounts to a total of  $O(|E(U_{ab})| \log n)$ .

We also have to determine all new degrees and the new excess numbers. This concerns all vertices of  $U_{ab}$ . Every such vertex is contained in at most  $2 \log n$  graphs  $U_{xy}^H$ . Hence all these numbers can be computed in  $O(\log n |U_{ab}|)$  time if all vertices of  $U_{ab}$  are removed. In other words, the data structure of  $H = G \setminus U_{ab}$  can be determined from that of  $G$  in  $O(\log n |U_{ab}|)$  time, including all degrees, excess numbers etc. (In the course of the action we take note of the first peripheral  $U$ -sets we encounter.)

We now repeat this process by removing peripheral  $U$ -sets until we reach  $K_1$ . The total complexity is then  $O(\log n \sum |U_{ab}|) = O(m \log n)$ .  $\square$

Now all prerequisites are ready for the following algorithm that recognizes whether a given graph  $G$  is a median graph with  $g(G) = 2$ .

**Algorithm 1**

- Input:* The adjacency list of a graph  $G$ .
- Output:* YES and a list of all geodetic pairs if  $G$  is a median graph with  $g(G) = 2$ .  
NO otherwise.
- Step 0:* If  $\Delta(G) > 2 \log_2 n$ , reject.  
If  $G$  is not a median graph, reject.  
Otherwise determine all  $\Theta$ -classes and the adjacency lists of all  $U$ -sets.  
Set  $i = k$ , where  $k = \text{idim}(G)$ , and  $G_k = G$ .

- Step 1: Compute the excess for all vertices in the  $U$ -sets and of all  $U$ -sets.*  
*Step 2: Find a peripheral  $U_{ab}$  as in Theorem 4.8.*  
*Step 3: Remove  $U_{ab}$  to obtain  $G_{i-1}$ .*  
*Step 4: Repeat step 2 and 3 (sequence of contractions) until  $G_0 = K_1$ .*  
*Step 5: For  $i = 0$  to  $k - 1$  do:*  
     *Find all geodetic pairs of  $G_i$  and determine those of  $G_{i+1}$  with the aid of Corollary 4.7.*  
*Step 6: If there are no such sets, return NO.*  
     *Otherwise return YES and the list of all geodetic pairs.*

**Theorem 4.9** *Let  $G$  be a graph  $G$ . Then Algorithm 1 correctly recognizes whether  $G$  is a median graph with  $g(G) = 2$ . It can be implemented to run in  $O(m \log n)$  time.*

**Proof.** Combining Lemma 4.1 and Proposition 4.4 we infer that Step 0 can be implemented in  $O(m \log n)$  time. Steps 1–4 are an algorithmic interpretation of the proof of Theorem 4.8. As stated in Theorem 4.8, one can perform these steps in  $O(m \log n)$  time. From Corollary 4.7 we find that Step 5 can also be performed in the desired time.  $\square$

## 5 Remoteness function in hypercubes

In this section we study the remoteness function in hypercubes which form the fundamental example of median graphs. Some of the results will be used in the last section where we will consider antimedians in median graphs in relation with their embeddings in hypercubes.

For a vertex  $x$  of  $Q_n$  let  $\bar{x}$  be its antipodal vertex, that is, the vertex that is obtained from  $x$  by reversing the roles of zeros and ones. Let  $X \subseteq V(Q_n)$ . Then

$$\bar{X} = \{\bar{x} \mid x \in X\}$$

is called the *antipodal set of  $X$* . Since  $\bar{\bar{x}} = x$  for  $x \neq \bar{x}$  it follows that  $\bar{\bar{X}} = X$ .

Let  $\pi = (x_1, \dots, x_k)$  be a profile on  $Q_d$ . For  $i = 1, \dots, k$  let  $n_0^{(i)}$  and  $n_1^{(i)}$  be the number of vertices from  $\pi$  with the  $i$ th coordinate equal 0 and 1, respectively. More formally,

$$n_0^{(i)}(\pi) = |\{x \in \pi \mid x^{(i)} = 0\}|$$

and

$$n_1^{(i)}(\pi) = |\{x \in \pi \mid x^{(i)} = 1\}|.$$

Define  $\text{Majority}(\pi)$  as the set of vertices  $u = u^{(1)} \dots u^{(d)}$  of  $Q_d$ , where

$$u^{(i)} \begin{cases} = 0; & n_0^{(i)}(\pi) > n_1^{(i)}(\pi), \\ = 1; & n_0^{(i)}(\pi) < n_1^{(i)}(\pi), \\ \in \{0, 1\}; & n_0^{(i)}(\pi) = n_1^{(i)}(\pi). \end{cases}$$

We say that vertices  $u \in \text{Majority}(\pi)$  are obtained by the *majority rule*.  $\text{Minority}(\pi)$  and the *minority rule* are defined analogously. It is easy to verify (using that the distance between vertices in hypercubes coincides with their Hamming distance) that  $M(\pi, Q_n) = \text{Majority}(\pi)$ , and similarly  $AM(\pi, Q_n) = \text{Minority}(\pi)$ . We now infer:

**Lemma 5.1** *Let  $\pi$  be a profile on  $Q_n$ . Then  $M(\pi, Q_n)$  induces a subcube of  $Q_n$ . Moreover,  $AM(\pi, Q_n) = M(\pi, Q_n)$ .*

Let  $Q$  and  $Q'$  be two subcubes of  $Q_n$ . Then we say that  $Q$  and  $Q'$  are *parallel* if they are of the same dimension, say  $r$ , and if vertices  $v_i$  of  $Q$  and  $v'_i$  of  $Q'$  can be ordered such that  $d(v_i, v'_i) = s$  for some integer  $s$  and for any  $i = 1, 2, \dots, 2^r$ , where the mapping  $v_i \mapsto v'_i$  is an isomorphism  $Q \rightarrow Q'$ .

**Proposition 5.2** *Let  $\pi$  be a profile on  $Q_n$  and let  $Q$  be a subcube parallel to the subcube induced by  $M(\pi, Q_n)$ . Then the function  $D(\cdot, \pi)$  is constant on  $Q$ .*

**Proof.** If  $|M(\pi, Q_n)| = 1$  there is nothing to be proved. Assume in the rest that  $|M(\pi, Q_n)| > 1$ , hence  $|\pi|$  must be even. By Lemma 5.1,  $M(\pi, Q_n)$  induces a subcube  $Q'$  and let  $x'y'$  be an edge of  $Q'$ . Partition the profile  $\pi$  into subprofiles  $\pi_1$  and  $\pi_2$ , where vertices of  $\pi_1$  lie in  $W_{x'y'}$  and vertices of  $\pi_2$  in  $W_{y'x'}$ . Since  $x', y' \in M(\pi, Q_n)$ , we have  $D(x', \pi) = D(y', \pi)$ . Therefore, the following reasoning

$$\begin{aligned} D(x', \pi) &= D(x', \pi_1) + D(x', \pi_2) \\ &= D(y', \pi_1) - |\pi_1| + D(y', \pi_2) + |\pi_2| \\ &= D(y', \pi_1) + D(y', \pi_2) = D(y', \pi) \end{aligned}$$

implies that  $|\pi_1| = |\pi_2|$ .

Let  $d(Q, Q') = s$  and let  $xy$  be the edge of  $Q$  with  $d(x, x') = d(y, y') = s$ . Then  $xy \Theta x'y'$  and consequently  $W_{xy} = W_{x'y'}$  and  $W_{yx} = W_{y'x'}$ . From the definition of  $W_{xy}$  and because  $|\pi_1| = |\pi_2|$  it follows that  $D(x, \pi) = D(y, \pi)$ . By the connectivity of  $Q$  we conclude that  $D$  must be a constant function on  $Q$ .  $\square$

We can generalize the concept of antipodes from hypercubes to Hamming graphs, noting that an antipode of a vertex  $x$  is any vertex that is farthest from  $x$ . In the case of hypercubes this vertex is unique, but not in general Hamming graphs. Hence for a vertex  $x$  of a Hamming graph  $H$  its *antipodal vertex* is any vertex  $y$  such that  $y^{(i)} \neq x^{(i)}$  for all  $i = 1, \dots, m$ . For  $X \subseteq V(H)$ , let the *antipodal set*  $\bar{X}$  of  $X$  be the set of all antipodal vertices over all vertices of  $X$ .

**Theorem 5.3** *A Hamming graph  $H$  is a hypercube if and only if for any profile  $\pi$*

$$AM(\pi, H) = \overline{M(\pi, H)}.$$

**Proof.** Suppose  $H$  is a hypercube. Then  $AM(\pi, H) = \overline{M(\pi, H)}$  for any profile  $\pi$  by Lemma 5.1.

For the converse suppose that a Hamming graph  $H$  is not a hypercube and let  $j$  be the index (coordinate) with  $m_j \geq 3$ . Consider the following profile  $\pi = (x, y)$  of size 2 such that  $x^{(i)} = y^{(i)} = 0$  for all  $i \neq j$  and let  $x^{(j)} = 0, y^{(j)} = 1$ . Then  $M(\pi, H) = \{x, y\}$ , and  $\overline{M(\pi, H)}$  consists of vertices  $z$  with  $z^{(i)} > 0$  for  $i \neq j$ . On the other hand  $AM(\pi, H)$  consists of vertices  $z$  with  $z^{(i)} > 0$  for  $i \neq j$  and  $z^{(j)} > 1$ . Hence  $AM(\pi, H) \subset \overline{M(\pi, H)}$  and the inclusion is strict, by which the theorem is proved.  $\square$

## 6 The remoteness function in median graphs, embedded in a hypercube

In this section we obtain some properties of the remoteness function in arbitrary median graphs, by using their isometric embedding into hypercubes. Since the properties of median sets have already been studied in several papers, we restrict mainly to the properties of antimedian sets in median graphs.

A vertex  $v$  of  $G$  is called a *local minimum* of a function  $D(x, \pi)$  if  $D(v, \pi) \leq D(u, \pi)$  for any neighbor  $u$  of  $v$ . It was proved in [5] that in a graph  $G$  the set  $M(\pi, G)$  is connected for any profile  $\pi$  on  $G$  if and only if for any  $\pi$  the function  $D(x, \pi)$  has the property that every local minimum is a global minimum. Since median graphs have the property that  $M(\pi, G)$  is connected for every  $\pi$ , we derive that in median graphs every local minimum is a global minimum.

For antimedian vertices (in other words, vertices achieving global maximum of  $D(x, \pi)$ ) the analogous result is not true for median graphs. Consider for example the  $3 \times 4$  grid, and one of the two vertices of degree 4 as the only vertex of the profile  $\pi$  (all four vertices of degree 2 achieve a local maximum, but only two of them are also global). Thus there are local maxima which are not global maxima and, moreover, antimedians need not be connected.

Restricting to hypercubes the fact that local minima are global minima can be strengthened as follows. First recall that by Lemma 5.1, the median of  $\pi$  is a subcube in  $Q_n$ , and the antimedian is its antipodal (hence parallel) subcube. By Proposition 5.2,  $D(x, \pi)$  is constant on every subcube parallel to them. Hence on any two shortest paths from  $M(\pi, Q_n)$  to  $AM(\pi, Q_n)$ , the two corresponding sequences of values of the remoteness function are the same. (Note also that any two distinct intervals from vertices in  $M(\pi, Q_n)$  to their (unique) closest vertices in  $AM(\pi, Q_n)$

are disjoint, and every vertex of  $G$  lies on some shortest path from  $M(\pi, Q_n)$  to  $AM(\pi, Q_n)$ .)

**Lemma 6.1** *Let  $\pi$  be a profile on  $Q_n$  and let  $xx'$  be an edge of  $Q_n$  such that  $d(x', AM(\pi, Q_n)) < d(x, AM(\pi, Q_n))$ . Then  $D(x, \pi) < D(x', \pi)$ .*

**Proof.** Let  $k = |\pi|$  and let  $m_j = \min\{n_0^{(j)}(\pi), n_1^{(j)}(\pi)\}$  and  $M_j = \max\{n_0^{(j)}(\pi), n_1^{(j)}(\pi)\}$ . Since  $AM(\pi, Q_n)$  can be obtained by the minority rule, for all  $a \in AM(\pi, Q_n)$ , we have

$$D(a, \pi) = \sum_{j=1}^n M_j.$$

Let  $d(x', AM(\pi, Q_n)) = d(x', a_{x'}) = l$ , where  $a_{x'}$  is the unique closest vertex to  $x'$  from  $AM(\pi, Q_n)$ . Then

$$\begin{aligned} D(x', \pi) &= D(a_{x'}, \pi) - \sum_{p=1}^l M_{i_p} + \sum_{p=1}^l m_{i_p}, \\ &= D(a_{x'}, \pi) - \sum_{p=1}^l (M_{i_p} - m_{i_p}) \end{aligned}$$

where  $x'$  and  $a_{x'}$  differ at coordinates  $i_p$ ,  $p = 1, \dots, l$ . Since  $x, x'$  are adjacent and  $d(x, a_{x'}) = d(x', a_{x'}) + 1$  there exists a coordinate  $p_{l+1}$ , distinct from all coordinates  $i_p$ ,  $1 \leq p \leq l$ , such that

$$D(x, \pi) = D(a_{x'}, \pi) - \sum_{p=1}^{l+1} (M_{i_p} - m_{i_p})$$

and  $D(x, \pi) < D(x', \pi)$ . □

**Theorem 6.2** *Let  $G$  be a median graph embedded isometrically into  $Q_n$ , and let  $\pi$  be a profile on  $G$ . Let  $a \in AM(\pi, G)$  and let  $a'$  be the closest vertex to  $a$  in  $AM(\pi, Q_n)$ . Then*

$$I(a, a') \cap V(G) = \{a\}.$$

**Proof.** Let  $b$  be the closest vertex to  $a'$  in  $M(\pi, Q_n)$ . From Lemma 5.1 we find that  $b$  is unique (as subcubes of a cube are gated; see [13], if necessary). In addition, Lemma 6.1 implies that  $D(x, \pi)$  is strictly increasing on any shortest path from  $b$  to  $a'$ . Since  $I(a, a') \subseteq I(b, a')$ , it follows that  $D(x, \pi)$  is strictly increasing on any shortest path from  $a$  to  $a'$ . Thus  $c \in I(a, a') \cap V(G)$ ,  $c \neq a$ , would imply that  $D(c, \pi) > D(a, \pi)$ , a contradiction with  $a \in AM(\pi, G)$ . Hence  $I(a, a') \cap V(G) = \{a\}$ . □




Figure 2: Example on antimedian.

In Fig. 2 we give an illustration of the above theorem. Vertices of a median graph  $G$  are darkened, and  $G$  is isometrically embedded into the 3-cube. Let the profile  $\pi$  consist of all five vertices of  $G$ . Then  $AM(\pi, Q_3)$  consists of the vertex  $w$ , where  $D(w, \pi) = 10$ . Vertices  $u$  and  $v$  are the only vertices from  $G$  that enjoy the condition from the theorem, that is  $I(a, w) \cap V(G) = \{a\}$ . Hence  $u$  and  $v$  are the only candidates to be antimedian vertices with respect to  $G$ , and both achieve the local maximum of  $D(\cdot, \pi)$  with respect to  $G$ . Since  $D(u, \pi) = 8$  and  $D(v, \pi) = 7$ , we infer that  $AM(\pi, G) = \{u\}$ . Note that even though  $v$  is closer to  $AM(\pi, Q_n)$  (that is, to  $w$ ) than  $u$ , it is not an antimedian vertex.

We proved in [3] that  $M(\pi, Q_n) \cap V(G) \neq \emptyset$  holds for any profile  $\pi$  which is used in an efficient algorithm for computing median sets in median graphs. In the events when  $AM(\pi, Q_n) \cap V(G) \neq \emptyset$  we have  $AM(\pi, G) = AM(\pi, Q_n) \cap V(G)$ , and then the antimedian set is also connected and it induces isometric subgraph of  $G$ . Unfortunately  $AM(\pi, Q_n) \cap V(G) \neq \emptyset$  is not true in general, as can be seen in the example from Fig. 2. Nevertheless, Theorem 6.2 could occasionally be helpful in finding the antimedian set for profiles on median graphs, since it can considerably reduce the number of candidates for the antimedian set to the vertices that achieve the condition from the theorem.

## References

- [1] A.V. Aho, J.E. Hopcroft and J.D. Ullman, *The Design and Analysis of Computer Algorithms*, Addison-Wesley, Reading, MA 1974.
- [2] K. Balakrishnan, *Algorithms for Median Computation in Median Graphs and their Generalizations Using Consensus Strategies*, Ph.D Thesis, University of Kerala, 2006.
- [3] K. Balakrishnan, B. Brešar, M. Changat, S. Klavžar, M. Kovše and A. R. Subhamathi, *Computing median and antimedian sets in median graphs*, submitted.

- [4] H.-J. Bandelt and J.-P. Barthélemy, Medians in median graphs, *Discrete Appl. Math.* 8 (1984) 131–142.
- [5] H.-J. Bandelt and V. Chepoi, Graphs with connected medians, *SIAM J. Discrete Math.* 15 (2002) 268–282.
- [6] B. Brešar and A. Tepoh Horvat, On the geodetic number of median graphs, *Discrete Math.* in press.
- [7] P. Cappanera, G. Gallo and F. Maffioli, Discrete facility location and routing of obnoxious activities, *Discrete Appl. Math.* 133 (2003) 3–28.
- [8] D. Djoković, Distance preserving subgraphs of hypercubes, *J. Combin. Theory Ser. B* 14 (1973) 263–267.
- [9] J. Hagauer, W. Imrich and S. Klavžar, Recognizing median graphs in sub-quadratic time, *Theoret. Comput. Sci.* 215 (1999) 123–136.
- [10] W. Imrich and S. Klavžar, Recognizing graphs of acyclic cubical complexes, *Discrete Appl. Math.* 1999 (95) 321–330.
- [11] W. Imrich and S. Klavžar, *Product Graphs: Structure and Recognition*, Wiley, New York, 2000.
- [12] W. Imrich, S. Klavžar and H. M. Mulder, Median graphs and triangle-free graphs, *SIAM J. Discrete Math.* 12 (1999) 111–118.
- [13] S. Klavžar and H. M. Mulder, Median graphs: characterizations, location theory and related structures, *J. Combin. Math. Combin. Comput.* 30 (1999) 103–127.
- [14] B. Leclerc, The median procedure in the semilattice of orders, *Discrete Appl. Math.* 127 (2003) 285–302.
- [15] F. R. McMorris, H. M. Mulder and F. R. Roberts, The median procedure on median graphs, *Discrete Appl. Math.* 84 (1998) 165–181.
- [16] H. M. Mulder, Majority strategy on graphs, *Discrete Appl. Math.* 80 (1997) 97–105.
- [17] H. M. Mulder, *The Interval Function of a Graph*, Math. Centre Tracts 132, Mathematisch Centrum, Amsterdam, 1980.
- [18] H. M. Mulder, *The expansion procedure for graphs*, in: R. Bodendiek ed., *Contemporary Methods in Graph Theory*, B.I.-Wissenschaftsverlag, Mannheim/Wien/Zürich (1990), 459–477.
- [19] S. B. Rao and A. Vijaykumar, On the median and the antimedial of a cograph, manuscript.

- [20] A. Tamir, Locating two obnoxious facilities using the weighted maximin criterion, *Oper. Res. Lett.* 34 (2006) 97–105.
- [21] P. Winkler, Isometric embeddings in products of complete graphs, *Discrete Appl. Math.* 7 (1984) 221–225.
- [22] B. Zmazek and J. Žerovnik, The obnoxious center problem on weighted cactus graphs, *Discrete Appl. Math.* 136 (2004) 377–386.