

UNIVERSITY OF LJUBLJANA  
INSTITUTE OF MATHEMATICS, PHYSICS AND MECHANICS  
DEPARTMENT OF MATHEMATICS  
JADRANSKA 19, 1000 LJUBLJANA, SLOVENIA

**Preprint series, Vol. 47 (2009), 1082**

A 2-PARAMETRIC  
GENERALIZATION OF  
SIERPIŃSKI GASKET GRAPHS

Marko Jakovac

ISSN 1318-4865

Ljubljana, March 2, 2009

# A 2-parametric generalization of Sierpiński gasket graphs

Marko Jakovac

Department of Mathematics and Computer Science

Faculty of Natural Sciences and Mathematics, University of Maribor

Koroška cesta 160, 2000 Maribor, Slovenia

marko.jakovac@uni-mb.si

February 25, 2009

## Abstract

Graphs  $S[n, k]$  are introduced as the graphs obtained from the Sierpiński graphs  $S(n, k)$  by contracting edges that lie in no triangle. The family  $S[n, k]$  is a previously studied class of Sierpiński gasket graphs  $S_n$ . Several properties of graphs  $S[n, k]$  are studied in particular, hamiltonicity and chromatic number.

**Key words:** Sierpiński graphs; Sierpiński gasket graphs; Hamiltonicity; Chromatic number

**AMS subject classification (2000):** 05C15, 05C45

## 1 Introduction

Sierpiński-like graphs appear in many different areas of graph theory, topology, probability ([7, 10]), psychology ([15]), etc. The special case  $S(n, 3)$  turns out to be only a step away of the famous Sierpiński gasket graphs  $S_n$ —the graphs obtained after a finite number of iterations that in the limit give the Sierpiński gasket, see [9]. This connection was introduced by Grundy, Scorer and Smith in [21] and later observed in [22].

Graphs  $S(n, 3)$  are also important for the Tower of Hanoi game since they are isomorphic to Hanoi graphs with  $n$  discs and 3 pegs. Metric properties, planarity, vertex and edge coloring were studied by now, see, for instance [1, 4, 5, 6, 13, 20]. Furthermore, in [11] it is proved that graphs  $S(n, 3)$  are uniquely 3-edge-colorable and have unique Hamiltonian cycles.

Graphs  $S(n, 3)$  can be generalized to Sierpiński graphs  $S(n, k)$ ,  $k \geq 3$ . The motivation came from topological studies of the Lipscomb's space [16, 17]. Graphs  $S(n, k)$  independently appeared in [19]. These graphs have many interesting properties, for instance, coding [3] and metric properties [18]. Moreover, in [12] it is

shown that graphs  $S(n, k)$  are Hamiltonian and that there are at most two shortest paths between any pair of their vertices. The length between any two vertices can be determined in  $O(n)$  time.

Sierpiński graphs are almost regular. In a natural way, two new families of regular Sierpiński-like graphs  $S^+(n, k)$  and  $S^{++}(n, k)$  were introduced in [14] and their crossing numbers determined (in terms of the crossing number of complete graphs).

Since Sierpiński gasket graphs  $S_n$  are important and are naturally derived from Sierpiński graphs  $S(n, 3)$ , we can apply the same construction for  $S(n, k)$ , where  $k \geq 3$ . We do this in section 2 and denote the new graphs  $S[n, k]$ . It is well known that graphs  $S(n, k)$ ,  $k \geq 3$  are Hamiltonian [12], as are graphs  $S_n$  [23]. We shall prove the same result for graphs  $S[n, k]$ . The chromatic number, chromatic index and total chromatic number of the Sierpiński graphs  $S(n, k)$  were already determined [18, 8]. In [8], a question was posted for the total chromatic number, where  $k$  is even. The question was answered in [6]. The same chromatic properties were studied for the Sierpiński gasket graphs. The chromatic number was determined in [23], chromatic index in [11] and total chromatic number in [8]. In this paper, we study the chromatic number of graphs  $S[n, k]$ .

## 2 Graphs $S[n, k]$ and their basic properties

First we recall the definition of the Sierpiński graphs  $S(n, k)$ . They are defined for  $n \geq 1$  and  $k \geq 1$  as follows. The vertex set of  $S(n, k)$  consists of all  $n$ -tuples of integers  $1, 2, \dots, k$ , that is,  $V(S(n, k)) = \{1, 2, \dots, k\}^n$ . Two different vertices  $u = (u_1, \dots, u_n)$  and  $v = (v_1, \dots, v_n)$  are adjacent if and only if there exists an  $h \in \{1, \dots, n\}$  such that

- (i)  $u_t = v_t$ , for  $t = 1, \dots, h - 1$ ;
- (ii)  $u_h \neq v_h$ ; and
- (iii)  $u_t = v_h$  and  $v_t = u_h$  for  $t = h + 1, \dots, n$ .

We will shortly write  $u_1 u_2 \dots u_n$  for  $(u_1, u_2, \dots, u_n)$ . See Fig. 1 for  $S(2, 4)$ .

By fixing  $u_1 \in \{1, \dots, k\}$ , we get  $S(n - 1, k)$ . In other words,  $S(n, k)$  is constructed of  $k$  different  $S(n - 1, k)$ . We label each with  $S_i(n, k)$ , for every  $i \in \{1, \dots, k\}$ . Note that  $S_i(n, k)$  and  $S_j(n, k)$ ,  $i \neq j$ , are connected with a single edge between vertices  $ij \dots j$  and  $ji \dots i$ . As in [8] we call these edges the linking edges of  $S(n, k)$ .

In [11] in a natural way the Sierpiński gasket  $S_n$  is constructed from the Sierpiński graph  $S(n, 3)$ . We can apply the same construction method for any Sierpiński graph  $S(n, k)$ .

Let  $u_1 u_2 \dots u_r j l \dots l$  and  $u_1 u_2 \dots u_r l j \dots j$ ,  $0 \leq r \leq n - 2$ , be two adjacent vertices of graph  $S(n, k)$ . We identify them in one vertex and write  $u_1 u_2 \dots u_r \{j, l\}$  or shortly  $u_{(r)} \{j, l\}$ ,  $j \neq l$  and  $j, l \in \{1, \dots, k\}$ . We get a 2-parametric Sierpiński




Figure 1: Graph  $S(2, 4)$

gasket graph  $S[n, k]$  (shortly  $k$ -Sierpiński gasket graph). We already know the case for  $k = 3$ . It is simply called the Sierpiński gasket. An example of  $S[2, 4]$  is shown in the Fig. 2.

Since  $S(n, k)$  is built of  $k$  copies of  $S(n - 1, k)$ , the graph  $S[n, k]$  is also built of  $k$  copies of  $S[n - 1, k]$ . We denote each copy with  $S_i[n, k]$ . Note that  $S_i[n, k]$  and  $S_j[n, k]$ ,  $i \neq j$ , share one vertex, that is  $\{i, j\}$ .

It is known, that  $S(1, k)$  is isomorphic to  $K_k$ . Hence,  $S[1, k]$  is also isomorphic to  $K_k$ .

Adjacency of vertices in  $S[n, k]$  is given in the next proposition.

**Proposition 2.1** *Let  $n \geq 2$  and  $u = u_1 \dots u_r \{i, j\}$  be a vertex in  $S[n, k]$ ,  $i, j, l \in \{1, \dots, k\}$ ,  $i \neq j, j \neq l, i \neq l$ .*

(i) *If  $0 \leq r \leq n - 3$ , then  $u$  is adjacent to*

$$\bar{u}_{(n-2)}\{j, l\}, l \in \{1, \dots, k\} \setminus \{j\},$$

and

$$\bar{\bar{u}}_{(n-2)}\{i, l\}, l \in \{1, \dots, k\} \setminus \{i\}.$$

(ii) *If  $r = n - 2$ , then  $u$  is adjacent to*

$$\bar{u}_{(n-2)}\{i, l\}, l \in \{1, \dots, k\} \setminus \{i, j\},$$

$$\bar{\bar{u}}_{(n-2)}\{j, l\}, l \in \{1, \dots, k\} \setminus \{i, j\},$$




Figure 2:  $S[2, 4]$

$$\begin{cases} \bar{u}_{(t-1)}\{i, u_t\}, & t \text{ is the largest index such that } u_t \neq i, 1 \leq t \leq n-2, \\ i \dots i, & \text{else,} \end{cases}$$

and

$$\begin{cases} \bar{\bar{u}}_{(s-1)}\{j, u_s\}, & s \text{ is the largest index such that } u_s \neq j, 1 \leq s \leq n-2, \\ j \dots j, & \text{else.} \end{cases}$$

**Proof.** (i) Let  $r \leq n-3$ . Take two vertices  $\bar{u} = u_1 \dots u_r i j \dots j j$  and  $\bar{\bar{u}} = u_1 \dots u_r j i \dots i i$ , where  $\bar{u}$  ends with at least two  $j$ 's and  $\bar{\bar{u}}$  with at least two  $i$ 's. Then  $\bar{u}$  is in  $S(n, k)$  adjacent to  $u_1 \dots u_r i j \dots j l$ ,  $l \in \{1, \dots, k\} \setminus \{j\}$ . In the construction procedure this vertex contracts to  $\bar{u}_{n-2}\{j, l\}$ ,  $l \in \{1, \dots, k\} \setminus \{j\}$  in  $S[n, k]$ . Similarly,  $\bar{\bar{u}}$  is adjacent to  $u_1 \dots u_r j i \dots i l$ ,  $l \in \{1, \dots, k\} \setminus \{i\}$ , which contracts to  $\bar{\bar{u}}_{n-2}\{i, l\}$ ,  $l \in \{1, \dots, k\} \setminus \{i\}$  in  $S[n, k]$ . The argument also holds for  $r = 0$ .

(ii) Let  $r = n-2$ . Then  $\bar{u} = u_1 \dots u_{n-2} i j$  and  $\bar{\bar{u}} = u_1 \dots u_{n-2} j i$ . In  $S(n, k)$ , the vertex  $\bar{u}$  is adjacent to  $u_1 \dots u_{n-2} i l$ ,  $l \in \{1, \dots, k\} \setminus \{i, j\}$ , which contracts to  $\bar{u}_{n-2}\{i, l\}$ ,  $l \in \{1, \dots, k\} \setminus \{i, j\}$  in  $S[n, k]$ . It is also adjacent to  $x = u_1 \dots u_{n-2} i i$ . If  $u_1 = \dots = u_{n-2} = i$ , then  $\bar{u}$  is adjacent to the extreme vertex  $i \dots i$  in  $S[n, k]$ . If not all of  $u_1, \dots, u_{n-2}$  are  $i$ , then let  $t$  be the largest index that  $u_t \neq i$ . Then the vertex  $\bar{u}$  is adjacent to  $x = u_1 \dots u_t i \dots i$ ,  $t \in \{1, \dots, n-2\}$ . In this case,  $\bar{u}$  is adjacent to  $\bar{u}_{t-1}\{i, u_t\}$ . Similarly, the vertex  $\bar{\bar{u}}$  is adjacent to  $u_1 \dots u_{n-2} j l$ ,  $l \in \{1, \dots, k\} \setminus \{i, j\}$ ,

which contracts to  $\bar{u}_{n-2}\{i, l\}$ ,  $l \in \{1, \dots, k\} \setminus \{i, j\}$  in  $S[n, k]$ . The vertex  $\bar{u}$  is also adjacent to  $x = u_1 \dots u_{n-2} j j$ . If  $u_1 = \dots = u_{n-2} = j$ , then  $\bar{u}$  is adjacent to the extreme vertex  $j \dots j$  in  $S[n, k]$ . If not all of  $u_1, \dots, u_{n-2}$  are  $j$ , then let  $s$  be the largest index that  $u_s \neq j$ . Then the vertex  $\bar{u}$  is adjacent to  $y = u_1 \dots u_s j \dots j$ ,  $s \in \{1, \dots, n-2\}$ . Therefore,  $\bar{u}$  is adjacent to  $\bar{u}_{t-1}\{i, u_t\}$ .  $\square$

It is easy to see that the graph  $S[n, k]$  has  $k$  extreme vertices of degree  $k-1$  and  $|V(S[n, k])| - k$  remaining vertices of degree  $(k-1) + (k-1) = 2(k-1)$ . We immediately get

**Proposition 2.2** *Graph  $S[n, k]$  is Eulerian if and only if  $k$  is odd.*

To gain more knowledge of the structure of  $S[n, k]$ , we give the number of vertices and edges of  $S[n, k]$ .

**Proposition 2.3** *Graph  $S[n, k]$  has  $\frac{k}{2}(k^{n-1} + 1)$  vertices and  $k^{n-1} \binom{k}{2}$  edges.*

**Proof.** A vertex in  $S[n, k]$  is of the form  $u_1 u_2 \dots u_r \{j, l\}$ . We have  $k$  possibilities for every  $u_i$ ,  $i \in \{1, \dots, r\}$ , and  $\binom{k}{2}$  possibilities for the unordered pair  $\{j, l\}$ . There are also  $k$  extreme vertices left. We get:

$$|V(S[n, k])| = k + \sum_{r=0}^{n-2} k^r \cdot \binom{k}{2} = \frac{k}{2}(k^{n-1} + 1).$$

The graph  $S[n+1, k]$  has as much edges as  $k$  copies of  $S[n, k]$ . Therefore,

$$|E(S[n+1, k])| = k \cdot |E(S[n, k])| = k^n \cdot \binom{k}{2}.$$

$\square$

Applying Proposition 2.3 and setting  $k = 3$  we get:

**Corollary 2.4** *Graph  $S_n$  has  $\frac{3}{2}(3^{n-1} + 1)$  vertices and  $3^n$  edges.*

### 3 Hamiltonicity

In [23], Tequia and Godbole proved that graphs  $S_n$  are Hamiltonian. In this section we generalize their statement to  $S[n, k]$ . First, we need a lemma.

**Lemma 3.1** *The graph  $S[n, k]$ ,  $k \geq 2$ , has a Hamiltonian path connecting two arbitrary extreme vertices.*

**Proof.** The statement holds for  $k = 2$  because  $S[n, 2]$  is isomorphic to a path on  $2^{n-1} + 1$  vertices.

Let  $k \geq 3$ . This statement is true for  $n = 1$ , since  $S[1, k]$  is isomorphic to  $K_k$ . Let  $n \geq 2$ . Without loss of generality start in vertex  $1 \dots 1$ . Since  $S_1[n, k]$  is isomorphic to  $S[n - 1, k]$  we can find a Hamiltonian path from  $1 \dots 1$  to the vertex  $\{1, 2\}$ . With the same argument, we can find a Hamiltonian path in  $S_2[n, k]$  from the vertex  $\{1, 2\}$  to the vertex  $\{2, 3\}$ . Next we find a Hamiltonian path in  $S_3[n, k]$  from the vertex  $\{2, 3\}$  to the vertex  $\{3, 4\}$  by avoiding  $\{1, 3\}$  (since we locally have an induced complete graph, avoiding is possible). We continue the procedure in  $S_4[n, k]$  by finding a Hamiltonian path between vertices  $\{3, 4\}$  and  $\{4, 5\}$  by avoiding vertices  $\{1, 4\}$  and  $\{2, 4\}$ . In general we find a Hamiltonian path in  $S_i[n, k]$ ,  $i \in \{3, \dots, k-1\}$ , from the vertex  $\{i-1, i\}$  to the vertex  $\{i, i+1\}$  by avoiding vertices  $\{1, i\}, \{2, i\}, \dots, \{i-2, i\}$ . Finally, we find a Hamiltonian path in  $S_k[n, k]$  from vertex  $\{k-1, k\}$  to vertex  $k \dots k$  by avoiding vertices  $\{1, k\}, \{2, k\}, \dots, \{k-2, k\}$ . All together we have constructed a Hamiltonian path between vertices  $1 \dots 1$  and  $k \dots k$ .

Similarly, we can find a Hamiltonian path between any two different extreme vertices in  $S[n, k]$ .  $\square$




Figure 3: A Hamiltonian cycle in  $S[2, 4]$

**Theorem 3.2** *Graph  $S[n, k]$  is Hamiltonian, for any  $n \geq 1$  and  $k \geq 3$ .*

**Proof.** The statement is true for  $n = 1$ , since  $S[1, k]$  is a complete graph.

Let  $n \geq 2$ . By Lemma 3.1, we can find a Hamiltonian path from the vertex  $\{k, 1\}$  to the vertex  $\{1, 2\}$  in  $S_1[n, k]$ . Similarly, we can find a Hamiltonian path between

vertices  $\{1, 2\}$  and  $\{2, 3\}$  in  $S_2[n, k]$ . Now we find a Hamiltonian path in  $S_3[n, k]$  from between vertices  $\{2, 3\}$  and  $\{3, 4\}$  by avoiding vertex  $\{1, 3\}$ . In general, we find a Hamiltonian path in  $S_i[n, k]$  between vertices  $\{i-1, i\}$  and  $\{i, i+1\}$ ,  $i \in \{3, \dots, k-1\}$  by avoiding vertices  $\{1, i\}, \{2, i\}, \dots, \{i-2, i\}$ . Finally, we find a Hamiltonian path in  $S_k[n, k]$  from the vertex  $\{k-1, k\}$  to the vertex  $\{k, 1\}$  by avoiding vertices  $\{2, k\}, \dots, \{k-2, k\}$ . Again, all the avoiding is possible because locally we have an induced complete graph. All the paths together form a Hamiltonian cycle in  $S[n, k]$ .  $\square$

Fig. 3 shows a Hamiltonian cycle in  $S[2, 4]$  obtained with the above construction. Once again, by setting  $k = 3$  we get:

**Corollary 3.3** [23] *Graph  $S_n$  is Hamiltonian.*

## 4 On the chromatic number

Since  $S[n, k]$  is built of complete graphs  $K_k$ , it is obvious that  $\chi(S[n, k]) \geq k$ . We prove the following.

**Theorem 4.1** *For any  $n \geq 1$  and any  $k \geq 1$ ,  $\chi(S[n, k]) = k$ .*

**Proof.** For  $k = 1$  and  $k = 2$  we get a vertex and a path on  $2^{n-1} + 1$  vertices respectively for which the statement is true. Let  $k \geq 3$ . For  $n = 1$  we have a complete graph on  $k$  vertices. It is well known that  $\chi(K_k) = k$ . Let  $n \geq 2$ .

Graph  $S[n, k]$  consists of  $k$  copies of  $S[n-1, k]$ , each denoted with  $S_i[n, k]$ ,  $i \in \{1, \dots, k\}$ . Two copies, say  $S_i[n, k]$  and  $S_j[n, k]$ ,  $i \neq j$ , share a common vertex  $\{i, j\}$ . For every  $i, j \in \{1, \dots, k\}$ ,  $i \neq j$ , expand this vertex into two vertices  $ij \dots j$  and  $ji \dots i$  connected with a linking edge

Note that for  $n \geq 3$  the expansion process described above is not the inverse process of getting  $S[n, k]$  from the  $S(n, k)$ , since we do not expand all the vertices that were contracted in  $S(n, k)$ .

By induction assumption graph  $S_1[n, k]$  can be colored with  $k$  colors. Denote the color of the vertex  $1j \dots j$ ,  $j \in \{1, \dots, k\}$ , with  $c_{1+j-2 \pmod k}$ , where  $c_{i+j-2 \pmod k} \in \{1, \dots, k\}$ . Then graphs  $S_i[n, k]$ ,  $i \in \{2, \dots, k\}$ , can also be colored in such a way that a vertex  $ij \dots j$  receives color  $c_{i+j-2 \pmod k}$ . This coloring is the same then the coloring used to color graph  $S_1[n, k]$ , only to be rotated clockwise. In other words we color graph  $S[n-1, k]$  like  $S_1[n, k]$  and rotate it clockwise  $i-1$ -times to get the coloring of graph  $S_i[n, k]$ . See Fig. 4 for the visualization of the rotated colorings.

A quick observation is that we obviously do not get a proper vertex coloring of the expanded graph  $S[n, k]$  since vertices  $ij \dots j$  and  $ji \dots i$ ,  $i \neq j$ , receive the same color, that is  $c_{i+j-2 \pmod k}$ . By contracting the previously expanded linking edges,






Figure 4: Vertex coloring of graph  $S[n, k]$

and getting  $S[n, k]$ , the merged vertex  $\{i, j\}$  receives color  $c_{i+j-2 \pmod k}$ . Therefore, we get a proper vertex coloring of  $S[n, k]$ .  $\square$

By setting  $k = 3$  we immediately get the next result.

**Corollary 4.2** [23] *For any  $n \geq 1$ ,  $\chi(S_n) = 3$ .*

We conclude the paper by asking what is the chromatic index and the total chromatic number of  $S[n, k]$ ?

## Acknowledgment

I would like to thank Sandi Klavžar for many useful comments and for turning my attention to the new family of graphs.

## References

- [1] D. Berend and A. Sapir, The diameter of Hanoi graphs, *Inform. Process. Lett.* 98 (2006) 79–85.
- [2] C. N. Campos and C. P. de Mello, A result on the total colouring of powers of cycles, *Discrete Appl. Math.* 155 (2007) 585–597.
- [3] S. Gravier, S. Klavžar and M. Mollard, Codes and  $L(2, 1)$ -labelings in Sierpiński graphs, *Taiwan. J. Math.* 9 (2005) 671–681.
- [4] A. Hinz, S. Klavžar, U. Milutinović, D. Parisse and C. Petr, Metric properties of the Tower of Hanoi graphs and Stern’s diatomic sequence, *European J. Combin.* 26 (2005) 693–708.

- [5] A. M. Hinz and D. Parisse, On the planarity of Hanoi graphs, *Expo. Math.* 20 (2002) 263–268.
- [6] A. M. Hinz and D. Parisse, Coloring Hanoi and Sierpiński graphs, manuscript, 2006.
- [7] A. M. Hinz and A. Schief, The average distance on the Sierpiński gasket, *Probab. Theory Related Fields* 87 (1990) 129–138.
- [8] M. Jakovac and S. Klavžar, Vertex-, edge-, and total-colorings of Sierpiński-like graphs, *Discrete Math.*, in press. doi:10.1016/j.disc.2008.02.026.
- [9] A. Jonsson, A trace theorem for the Dirichlet form on the Sierpinski gasket, *Math. Z.* 250 (2005) 599–609.
- [10] V. A. Kaimanovich, Random walks on Sierpiński graphs: hyperbolicity and stochastic homogenization, *Fractals in Graz 2001*, 145–183, Trends Math., Birkhäuser, Basel, 2003.
- [11] S. Klavžar, Coloring Sierpiński graphs and Sierpiński gasket graphs, *Taiwan. J. Math.* 12 (2008) 513–522
- [12] S. Klavžar and U. Milutinović, Graphs  $S(n, k)$  and a variant of the Tower of Hanoi problem, *Czechoslovak Math. J.* 47(122) (1997) 95–104.
- [13] S. Klavžar, U. Milutinović and C. Petr, Hanoi graphs and some classical numbers, *Expo. Math.* 23 (2005) 371–378.
- [14] S. Klavžar and B. Mohar, Crossing numbers of Sierpiński-like graphs, *J. Graph Theory* 50 (2005) 186–198.
- [15] F. Klix and K. Rautenstrauch-Goede, Struktur- and Komponentenanalyse von Problemlösungsprozessen, *Z. Psychol.* 174 (1967) 167–193.
- [16] S. L. Lipscomb and J. C. Perry, Lipscomb’s  $L(A)$  space fractalized in Hilbert’s  $l^2(A)$  space, *Proc. Amer. Math. Soc.* 115 (1992) 1157–1165.
- [17] U. Milutinović, Completeness of the Lipscomb space, *Glas. Mat. Ser. III* 27(47) (1992) 343–364.
- [18] D. Parisse, On some metric properties of the Sierpiński graphs  $S(n, k)$ , *Ars Combin.* 90 (2009) 145–160.
- [19] T. Pisanski and T. W. Tucker, Growth in repeated truncations of maps, *Atti Sem. Mat. Fis. Univ. Modena* 49 (2001) 167–176.
- [20] D. Romik, Shortest paths in the Tower of Hanoi graph and finite automata, *SIAM J. Discrete Math.* 20 (2006) 610–622.

- [21] R. S. Scorer, P. M. Grundy and C. A. B. Smith, Some binary games, *Math. Gaz.* 28 (1944) 96–103.
- [22] H. Sydow, Zur metrischen Erfassung von subjektiven Problemzuständen und zu deren Veränderung im Denkprozeß, *Z. Psychol.* 177 (1970) 145–198.
- [23] A. M. Teguaia and A. P. Godbole, Sierpiński gasket graphs and some of their properties, *Australas. J. Combin.* 35 (2006) 181–192.
- [24] D. B. West, *Introduction to Graph Theory*, Second Edition, Prentice Hall, Inc., Upper Saddle River, NJ, 2001.